

The generator conjecture for 3^G subfactor planar algebras

Zhengwei Liu and David Penneys

February 15, 2017

Abstract


We state a conjecture for the formulas of the depth 4 low-weight rotational eigenvectors and their corresponding eigenvalues for the 3^G subfactor planar algebras. We prove the conjecture in the case when $|G|$ is odd. To do so, we find an action of G on the reduced subfactor planar algebra at $f^{(2)}$, which is obtained from shading the planar algebra of the even half. We also show that this reduced subfactor planar algebra is a Yang-Baxter planar algebra.

This is the published version of [arXiv:1507.04794](https://arxiv.org/abs/1507.04794).

Dedicated to the 60th birthday of Vaughan F. R. Jones.


1 Introduction

Haagerup initiated the classification of subfactor principal graphs with index a little greater than 4, and he gave a classification of all possible graph pairs in the index range $(4, 3 + \sqrt{2})$ [Haa94]. In doing so, he discovered a so-called ‘exotic’ subfactor [AH99] with index $\frac{5+\sqrt{13}}{2}$ and principal graph the 3-spoke




The $\mathbb{Z}/3\mathbb{Z}$ -symmetry of this graph means that the dimension one vertices at the ends of the spokes form the group $\mathbb{Z}/3\mathbb{Z}$ under the fusion operation of the corresponding bimodules.

In [Izu01], Izumi gave a generalized construction to other abelian groups using Cuntz algebras, and constructed an example when $G = \mathbb{Z}/5\mathbb{Z}$. Such a subfactor is called a 3^G subfactor and has principal graph



Two recent articles of Evans-Gannon [EG11, EG14] have successfully used Izumi’s equations to construct a myriad of new examples of 3^G and related $2^G 1$ subfactors with principal graphs



They also give simple formulas for the quantum double and its modular data, leading them to conjecture that there should be associated rational conformal field theories.

The even halves of the 3^G and $2^G 1$ subfactors are examples of quadratic unitary fusion categories, which have a group G of invertible objects and one other orbit $G\rho$ of simple objects, together with a relation for fusion on the right by g , and a quadratic fusion relation for ρ . For example, by unpublished work of Izumi, the even half of a 3^G subfactor for $|G|$ odd satisfies

$$\rho g = g^{-1} \rho \text{ and } \rho^2 \cong 1 \oplus \bigoplus_{g \in G} g\rho.$$

The even halves of $2^G 1$ subfactors are unitary near group fusion categories [EG14], which are generalizations of Tambara-Yamagami categories [TY98].

Izumi observed in [Izu01] that when realizing a unitary fusion category as a category of sectors of some infinite factor M , Cuntz algebras naturally appear as the C^* -algebras generated by the orthonormal bases of intertwiner spaces in M . Cuntz algebras are particularly useful in constructing quadratic categories because we usually only need to analyze one Cuntz algebra, in which the quadratic relation allows us to write down polynomial equations in the generators to define an endomorphism of the C^* -algebra. One then extends the endomorphisms to the von Neumann completion using the unique KMS state [OP78], which is again an infinite factor (e.g., see [BR97]). When the category is not quadratic, we obtain multiple Cuntz algebras together with relations between them, and the situation is much more complicated.

Currently planar algebra techniques are not as effective as Cuntz algebras for constructing quadratic categories. The recent articles [BP14, PP13] suggest a uniform skein theory for the 3^G 's using 2-strand jellyfish relations. A general formula for the generators in the graph planar algebras remains elusive, as the valence and size of the $2^n 1$ and 3^n graphs gets quite large. We expect that the Cuntz algebra and planar algebra techniques can be reconciled, which will be explored in future work. For example, Izumi has shown how to draw planar diagrams for the actions of his Cuntz algebra endomorphisms.

Based on [MP15, PP13] we conjecture specific formulas for the 3^G low-weight rotational eigenvectors in the 4-box spaces in terms of minimal projections in the 3^G subfactor planar algebras. This is the first step in the Jones-Peters graph planar algebra embedding program [Jon01, Pet10, JP11, Jon12] toward a uniform planar algebraic approach to the 3^G subfactors.

Let \mathcal{P}_\bullet be a 3^G subfactor planar algebra. For $g \in G \setminus \{1\}$, let p_g be the projection in $\mathcal{P}_{4,+}$ corresponding to $g\rho$. We make the following conjecture about the low-weight rotational eigenvectors for \mathcal{P}_\bullet which agrees with the Haagerup $3^{\mathbb{Z}/3\mathbb{Z}}$ and Izumi $3^{\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}}$ and $3^{\mathbb{Z}/4\mathbb{Z}}$ subfactor planar algebras by [Pet10, Jon12, MP15, PP13].


Conjecture A. *Suppose $g, h, k, \ell \in G \setminus \{1\}$ are distinct elements.*

- (1) *If $g = g^{-1}$ and $h = h^{-1}$, then $p_g - p_h$ is a low-weight rotational eigenvector with eigenvalue 1.*
- (2) *If $g = h^{-1}$, then $p_g - p_h$ is a low-weight rotational eigenvector with eigenvalue -1.*
- (3) *If $g = g^{-1}$ and $h = k^{-1}$, then $2p_g - (p_h + p_k)$ is a low-weight rotational eigenvector with eigenvalue 1.*
- (4) *If $g = h^{-1}$ and $k = \ell^{-1}$, then $(p_g + p_h) - (p_k + p_\ell)$ is a low-weight rotational eigenvector with eigenvalue 1.*


Our main theorem in this article is as follows.

Theorem B. *Conjecture A is true when $|G|$ is odd.*

To prove this theorem, we construct a G -action¹ on the reduced unshaded planar algebra \mathcal{R}_\bullet of \mathcal{P}_\bullet at $\rho = f^{(2)}$ with principal graphs



For each $g \in G$, we pick a distinguished isomorphism $V_g : \rho g^{-1} \rightarrow g\rho$. Denoting $\rho \in \mathcal{R}_1$ by a red strand and the group elements $g \in \mathcal{P}_{6,+}$ by black labelled, oriented strands, the action is given by



which is similar to diagrams arising from looking at connections [Ocn88, Jon99, MP14, Liu13]. Moreover, we have $\Phi_g \circ \Phi_h = \Phi_{gh}$, giving a G -action on \mathcal{R}_\bullet ¹. We anticipate this new technique will have new applications to subfactor planar algebras beyond the proof of our theorem.

When there is an $h \in G$ such that $h^2 = g$, we apply the action of Φ_h to the relation


$$\boxed{\begin{array}{|c|} \hline \text{---} \\ \hline \end{array}} = \boxed{\begin{array}{|c|} \hline \text{---} \\ \hline \end{array}} - \frac{1}{[3] - 1} \left(\boxed{\begin{array}{|c|} \hline \text{---} \\ \hline \end{array}} - \boxed{\begin{array}{|c|} \hline \text{---} \\ \hline \end{array}} \right),$$

where the trivalent vertex is a suitably normalized map $\rho \otimes \rho \rightarrow \rho$. We then obtain the formula

$$\mathcal{F}_{\mathcal{R}_\bullet}(p_g) = p_{g^{-1}} - \frac{1}{[3] - 1} \left(\boxed{\begin{array}{|c|} \hline \text{---} \\ \hline \end{array}} - \boxed{\begin{array}{|c|} \hline \text{---} \\ \hline \end{array}} \right).$$

By Corollary 3.9 below, Conjecture A is equivalent to this formula holding for every $g \in G$. When $|G|$ is odd, every $g \in G$ has a square root, but this is no longer the case when $|G|$ is even.

We note that the principal graph Λ above bears a strong resemblance to the principal graph of the reduced subfactor of A_7 at $f^{(2)}$, given by



It was recently shown in [LMP15] that the planar algebra corresponding to this reduced subfactor is a Yang-Baxter planar algebra, which is a planar algebra generated by 2-boxes with a relation which writes

¹There is actually a technicality involving the shading and the symmetric self-duality [MP15, MP] of \mathcal{R}_\bullet which we address in Sections 4 and 5, but we omit the shading in the introduction to give the spirit of the argument.

one type of triangle in terms of the other triangle and lower order terms (below, $\mathcal{B}_{2,+}$ is a basis of $\mathcal{P}_{2,+}$ including the diagrammatic basis of $\mathcal{TL}_{2,+}$):

$$\begin{array}{c} \star \\ \circ \\ a \\ \star \end{array} \begin{array}{c} \star \\ \circ \\ b \\ \star \end{array} = \sum_{x,y,z \in \mathcal{B}_{2,+}} \lambda_{x,y,z} \begin{array}{c} \star \\ \circ \\ x \\ \star \end{array} \begin{array}{c} \star \\ \circ \\ y \\ \star \end{array} \begin{array}{c} \star \\ \circ \\ z \\ \star \end{array} .$$

(There is also a relation swapping the above types of triangles, which is necessary to be able to evaluate all closed diagrams.) In his recent classification of singly generated Yang-Baxter planar algebras [Liu15], Liu discovered that the subfactor for \mathcal{S} belongs to an infinite family of subfactors arising from the E_{N+2} quantum subgroup of $SU(N)$.

We further conjecture a new skein theoretic approach to constructing the reduced subfactor planar algebra \mathcal{R}_\bullet of a 3^G subfactor planar algebra, and we prove it in the case $|G|$ is odd.

Conjecture C. \mathcal{R}_\bullet is a Yang-Baxter planar algebra with $|G| - 1$ generators.


Theorem D. Conjecture C is true when $|G|$ is odd.

1.1 Acknowledgements

The authors would like to thank Masaki Izumi and Emily Peters for helpful conversations; the Banff International Research Station for hosting the 2014 workshop on Subfactors and Fusion Categories, where this work started; and the 2014 Maui conference on Subfactor Theory in Mathematics and Physics, which was funded by NSF grant DMS-1400275 and DOD-DARPA grant HR0011-12-1-0009. Zhengwei Liu was supported by NSF grant DMS-1001560. David Penneys was partially supported by an AMS Simons travel grant and NSF grant DMS-1500387. Both authors were supported by DOD-DARPA grant HR0011-12-1-0009.

2 3^G subfactors

Definition 2.1. Let G be a non-trivial finite group. A 3^G subfactor planar algebra is a subfactor planar algebra whose principal graph Γ_+ is a $3^{|G|}$ spoke graph



where the even bimodules generate a G -quadratic category, denoted $\frac{1}{2}\mathcal{P}_+$, whose fusion rules are

- (1) $g \otimes h = gh$, i.e., we may identify the dimension 1 bimodules with G ,
- (2) $g \otimes \rho = g\rho$, so $\{g\rho | g \in G\}$ is a left G -set the obvious way: $g(h\rho) = (gh)\rho$,
- (3) $\rho \otimes g = \theta(g)\rho$ for some automorphism θ of G , since ρg is irreducible by Frobenius reciprocity, and
- (4) $\rho \otimes \rho = 1 \oplus \bigoplus_{g \in G} g\rho$.

Conjecture 2.2 (Izumi). *If a 3^G subfactor planar algebra exists, then G is abelian, and $\theta(g) = g^{-1}$ for all $g \in G$.*

Remark 2.3. In unpublished work, Izumi has proven Conjecture 2.2 for the case $|G|$ odd.

Assumption 2.4. We will assume G is abelian and $\theta(g) = g^{-1}$ for all $g \in G$.

Corollary 2.5. *Every bimodule at depth 4 is self-dual.*

Proof. For all $g \in G$, $\overline{g\rho} \cong \rho g^{-1} \cong \theta(g^{-1})\rho \cong g\rho$. □

An argument of Izumi gives the structure of the dual principal graph. We provide a proof for the reader’s convenience. We begin with a helpful lemma generalizing [MS12, Lemma 3.6]. For a pair of principal graphs (Γ_+, Γ_-) of a subfactor planar algebra \mathcal{P}_\bullet , let $\Gamma_\pm(n)$ denote the truncation to depth n . Denote the one-click rotation by \mathcal{F} .


Lemma 2.6. *Suppose Γ_\pm is exactly $(n - 1)$ supertransitive for an even $n \geq 2$. If $\Gamma_+(n)$ is simply-laced and only has self-dual vertices at depth n , then $\Gamma_-(n)$ is simply-laced and only has self-dual vertices at depth n .*

Proof. We analyze the rotation by π given by \mathcal{F}^n on $\mathcal{P}_{n,\pm} \ominus \mathcal{TL}_{n,\pm}$, where \mathcal{TL}_\bullet is the Temperley-Lieb planar subalgebra. Observe that the elements $p - \frac{\text{Tr}(p)}{[n+1]}f^{(n)}$ span $\mathcal{P}_{n,\pm} \ominus \mathcal{TL}_{n,\pm}$, where the p are the new projections at depth n . Since $\Gamma_+(n)$ is simply-laced and only has self-dual vertices, \mathcal{F}^n is the identity on $\mathcal{P}_{n,+} \ominus \mathcal{TL}_{n,+}$. Now \mathcal{F} and \mathcal{F}^* map Temperley-Lieb to Temperley-Lieb, so \mathcal{F} and \mathcal{F}^* also take the orthogonal complement to the orthogonal complement. Hence for all $x \in \mathcal{P}_{n,-} \ominus \mathcal{TL}_{n,-}$,


$$\mathcal{F}^n(x) = \mathcal{F}^n(\mathcal{F}(\mathcal{F}^{-1}(x))) = \mathcal{F}(\mathcal{F}^n(\mathcal{F}^{-1}(x))) = \mathcal{F}(\mathcal{F}^{-1}(x)) = x,$$

so \mathcal{F}^n is also the identity on $\mathcal{P}_{n,-} \ominus \mathcal{TL}_{n,-}$. Since the elements $p - \frac{\text{Tr}(p)}{[n+1]}f^{(n)}$ span $\mathcal{P}_{n,-} \ominus \mathcal{TL}_{n,-}$, where the p are the new projections at depth n of Γ_- , all the vertices at depth n of Γ_- are self dual. Now \mathcal{F}^n is also an anti-isomorphism of the algebra $\mathcal{P}_{n,-}$. Letting $\mathcal{I}_{n,-}$ be the basic construction ideal, we see that \mathcal{F}^n preserves $\mathcal{I}_{n,-}$, and passes to the quotient algebra $\mathcal{P}_{n,-}/\mathcal{I}_{n,-}$. But \mathcal{F}^n is the identity on $\mathcal{P}_{n,-}/\mathcal{I}_{n,-}$, and thus this algebra is abelian. Hence $\Gamma_-(n)$ is simply laced. □

Theorem 2.7 (Izumi). *For every self-dual tail of Γ_+ , there is self-dual tail of Γ_- .*



For every Haagerup tail of Γ_+ , there is a dual Haagerup tail of Γ_- .



Proof. We know the vertices of Γ_- at depth 3 and 5, so it remains to determine the vertices at depth 4 and 6 and the edges. First, we consider the vertices $\overline{X}g$ at depth 5. By Frobenius reciprocity,

$$\langle \overline{X}gX, \overline{X}gX \rangle = \langle X\overline{X}g, gX\overline{X} \rangle = \langle (1 + \rho)g, g(1 + \rho) \rangle = 1 + \langle \rho g, g\rho \rangle = \begin{cases} 2 & \text{if } g = g^{-1} \\ 1 & \text{if } g \neq g^{-1}. \end{cases}$$

Hence $\overline{X}gX$ is simple precisely when $g \neq g^{-1}$.

- Case 1: Suppose $g \neq g^{-1}$. Then $\overline{X}gX$ and $\overline{X}g^{-1}X$ are simple, and moreover,

$$\langle \overline{X}gX, \overline{X}g^{-1}X \rangle = \langle X\overline{X}g, g^{-1}X\overline{X} \rangle = \langle (1 + \rho)g, g^{-1}(1 + \rho) \rangle = 1,$$

so they are equal. Hence the distinct depth 5 vertices $\overline{X}g^{-1}$ and $\overline{X}g$ of Γ_- are univalent and connect to a single self-dual vertex $\overline{X}gX = \overline{X}g^{-1}X$ at depth 4. Since

$$\overline{X}gX\overline{X} = \overline{X}g(1 + \rho) = \overline{X}g + \overline{X}g\rho = \overline{X}g + \overline{X}\rho g^{-1} = \overline{X}g + \overline{X}g^{-1} + \overline{Z}g^{-1}, \quad (1)$$

each of which is simple, $\overline{X}gX$ connects by a single edge to the branch point at depth 3 of Γ_- .

- Case 2: Suppose $g = g^{-1}$. Then $\overline{X}g$ connects to two even vertices of Γ_- , at least one of which must connect to the branch point at depth 3. Since $\overline{X}gX\overline{X}$ splits into exactly three simples by Equation (1), $\overline{X}g$ must connect to one bivalent vertex at depth 4 and one univalent vertex at depth 6.

Now in both cases, $\overline{X}gX$ is self-dual, so the new vertices we have found so far at depths 4 and 6 are all self-dual, as the dual of a vertex must occur at the same depth. It remains to show there is a single self-dual vertex connected to \overline{Z} for each (unordered) set of elements $\{g, g^{-1}\}$ with $g \neq g^{-1}$.

Analyzing the Ocneanu 4-partite graph, we see there are $|G|$ paths from Z to \overline{Z} through $A - A$ bimodules, so there must be $|G|$ paths through $B - B$ bimodules. If $G = N \cup S \cup \{1\}$ where N is the set of non self-inverse elements of G and S is the set of non-trivial self-inverse elements of G , then currently we can account for $|S| + 1 + |N|/2$ paths through $B - B$ vertices.

Hence the remaining paths through $B - B$ bimodules must come from vertices at depth 4 which do not continue to depth 5. By Lemma 2.6, we know $\Gamma_-(4)$ is simply laced with only self-dual vertices, so there must be exactly $|N|/2$ self-dual univalent vertices at depth 4 of Γ_- . \square

For $g \in G \setminus \{1\}$, let p_g be the projection in $\mathcal{P}_{4,+}$ corresponding to $g\rho$. Let $\langle E_i \rangle$ denote the algebra generated by E_1, \dots, E_{n-1} in $TL_{n,+}$, and note that $\langle E_i \rangle$ is perpendicular to all projections on the principal graph and to the Jones-Wenzl idempotents, and $\text{span}(\{p_g | g \neq 1\}) = \mathcal{P}_{4,+} \ominus \langle E_i \rangle$.

Lemma 2.8. *If \mathcal{Q}_\bullet is an $n - 1$ supertransitive subfactor planar algebra, then any non-zero element in $\mathcal{Q}_{n,+} \ominus \langle E_i \rangle$ with zero trace is uncappable.*

Proof. Follows easily from $qE_i = 0$ for all $i < n$ and all minimal projections $q \in \mathcal{Q}_{n,+} \ominus \langle E_i \rangle$. \square

Corollary 2.9. *For all $g, h \in G \setminus \{1\}$ with $g \neq h$, $p_g - p_h$ is uncappable.*

Proposition 2.10. *The new low-weight vectors at depth 4 have eigenvalue ± 1 .*

Proof. By Corollary 2.5, \mathcal{F}^4 is the identity on $\text{span}(\{p_g | g \in G\})$. \square

2.1 The low weight rotational eigenvector conjecture

Conjecture (Conjecture A). *Suppose $g, h, k, \ell \in G \setminus \{1\}$ are distinct elements.*

- (1) *If $g = g^{-1}$ and $h = h^{-1}$, then $p_g - p_h$ is a low-weight rotational eigenvector with eigenvalue 1.*
- (2) *If $g = h^{-1}$, then $p_g - p_h$ is a low-weight rotational eigenvector with eigenvalue -1.*
- (3) *If $g = g^{-1}$ and $h = k^{-1}$, then $2p_g - (p_h + p_k)$ is a low-weight rotational eigenvector with eigenvalue 1.*

(4) If $g = h^{-1}$ and $k = \ell^{-1}$, then $(p_g + p_h) - (p_k + p_\ell)$ is a low-weight rotational eigenvector with eigenvalue 1.

Remark 2.11. Conjecture A agrees with the Haagerup $3^{\mathbb{Z}/3\mathbb{Z}}$ and Izumi $3^{\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}}$ and $3^{\mathbb{Z}/4\mathbb{Z}}$ subfactor planar algebras by [Pet10, Jon12, MP15, PP13]. See also Remark 2.14.

Lemma 2.12. The set $B = \{p_g | g \in G\} \cup \{\mathcal{F}^2(f^{(4)})\}$ is linearly independent.

Proof. First, note that $f^{(4)} = \sum_{g \in G \setminus \{1\}} p_g$ and $\mathcal{F}^2(f^{(4)})$ are linearly independent, since capping $\mathcal{F}^2(f^{(4)})$ on the bottom does not give zero. Suppose

$$0 = \lambda_f \mathcal{F}^2(f^{(4)}) + \sum_{g \in G \setminus \{1\}} \lambda_g p_g.$$

Taking inner products with p_g gives

$$0 = \lambda_g \text{Tr}(p_g) + \lambda_f \text{Tr}(\mathcal{F}^2(f^{(4)})p_g) = \lambda_g \text{Tr}(p_g) + \lambda_f \text{Tr}(p_g) (\text{coeff}_{\in f^{(4)}} \mathcal{F}^2(\text{id})),$$

so λ_g is independent of g . Call this constant λ . Then we have

$$0 = \lambda_f \mathcal{F}^2(f^{(4)}) + \lambda \sum_{g \in G \setminus \{1\}} p_g = \lambda_f \mathcal{F}^2(f^{(4)}) + \lambda f^{(4)},$$

so $\lambda_f = \lambda = 0$, and B is linearly independent. □

Proposition 2.13. Conjecture A holds if and only if for all $g \in G$,

$$\mathcal{F}^2(p_g) = \frac{1}{|G| - 1} (\mathcal{F}^2(f^{(4)}) - f^{(4)}) + p_{g^{-1}}.$$

Proof. If \mathcal{F}^2 is given by the above formula, a straightforward calculation shows that Conjecture A holds. We now prove the other direction.

Divide $G \setminus \{1\}$ into the two subsets: the non-trivial self-inverse elements S , and the non-self-inverse elements N . Let $N^+ \subset N$ so that for each $g \in N$, exactly one of $g, g^{-1} \in N^+$. Let $B_1 = \{p_g - p_{g^{-1}} | g \in N^+\}$, and note $|B_1| = |N|/2$.

Case 1: Suppose $S \neq \emptyset$, so $|G|$ is even. Fix $s_0 \in S$. Let $B_2 = \{2p_{s_0} - (p_g + p_{g^{-1}}) | g \in N^+\}$, and note $|B_2| = |N|/2$. Finally, let $B_3 = \{p_{s_0} - p_s | s \in S \setminus \{s_0\}\}$, and note $|B_3| = |S| - 2$. Observe $B' = B_1 \cup B_2 \cup B_3$ has size $|G| - 2$.

Claim. $D = B' \cup \{f^{(4)}, \mathcal{F}^2(f^{(4)})\}$ is a basis for $\text{span}(B)$.

Proof of Claim. It suffices to show D is linearly independent. Note that by taking linear combinations, we can obtain $p_g - p_h$ for all $g, h \in G \setminus \{1\}$. The result now follows since $f^{(4)} = \sum_{g \neq 1} p_g$. □

Now by Conjecture A,

$$[\mathcal{F}^2]_D = \begin{pmatrix} -I_{B_1} & 0 & 0 & 0 & 0 \\ 0 & I_{B_2} & 0 & 0 & 0 \\ 0 & 0 & I_{B_3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix},$$

and the change of basis matrix from D to B is given by

$$Q_D^B = \left(\begin{array}{c|cccccc} & B_1 = N^+ - N^- & B_2 = 2s_0 - N^+ & B_3 = s_0 - S \setminus \{s_0\} & f^{(4)} & \mathcal{F}^2(f^{(4)}) \\ \hline N^+ & I & -I & 0 & \mathbf{1} & 0 \\ N^- & -I & -I & 0 & \mathbf{1} & 0 \\ S \setminus \{s_0\} & 0 & 0 & -I & \mathbf{1} & 0 \\ s_0 & 0 & \mathbf{2} & \mathbf{1} & 1 & 0 \\ \mathcal{F}^2(f^{(4)}) & 0 & 0 & 0 & 0 & 1 \end{array} \right)$$

where I is an identity matrix, and \mathbf{k} is the matrix with all k 's.

Setting $n = |N^+| + |S| = |G| - 1$, it is straightforward to check

$$(Q_D^B)^{-1} = \frac{1}{2n} \left(\begin{array}{c|ccccc} & N^+ & N^- & S \setminus \{s_0\} & s_0 & \mathcal{F}^2(f^{(4)}) \\ \hline B_1 & nI & -nI & 0 & 0 & 0 \\ B_2 & \mathbf{2} - nI & \mathbf{2} - nI & \mathbf{2} & \mathbf{2} & 0 \\ B_3 & \mathbf{2} & \mathbf{2} & \mathbf{2} - 2nI & \mathbf{2} & 0 \\ f^{(4)} & \mathbf{2} & \mathbf{2} & \mathbf{2} & 2 & 0 \\ \mathcal{F}^2(f^{(4)}) & 0 & 0 & 0 & 0 & 2n \end{array} \right)$$

Hence

$$[\mathcal{F}^2]_B = Q_D^B [\mathcal{F}^2]_D (Q_D^B)^{-1} = \frac{1}{n} \left(\begin{array}{c|cccc} & N^+ & N^- & S & \mathcal{F}^2(f^{(4)}) \\ \hline N^+ & -\mathbf{1} & nI - \mathbf{1} & -\mathbf{1} & \mathbf{n} \\ N^- & nI - \mathbf{1} & -\mathbf{1} & -\mathbf{1} & \mathbf{n} \\ S & -\mathbf{1} & -\mathbf{1} & nI - \mathbf{1} & \mathbf{n} \\ \mathcal{F}^2(f^{(4)}) & \mathbf{1} & \mathbf{1} & \mathbf{1} & 0 \end{array} \right).$$

Case 2: Suppose $S = \emptyset$, so $|G|$ is odd. Fix $g_0 \in N^+$. Let

$$B_2 = \left\{ (p_g + p_{g^{-1}}) - (p_{g_0} + p_{g_0^{-1}}) \mid g \in N^+ \setminus \{g_0\} \right\},$$

and note $|B_2| = |N^+| - 1$. Hence if $D = B_1 \cup B_2$, we have $|D| = |G| - 2$. Similar to before, $D \cup \{f^{(4)}, \mathcal{F}^2(f^{(4)})\}$ is a basis for $\text{span}\{B\}$. Now by Conjecture A,

$$[\mathcal{F}^2]_D = \begin{pmatrix} -I_{B_1} & 0 & 0 & 0 \\ 0 & I_{B_2} & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix},$$

and the change of basis matrix from D to B is given by

$$Q_D^B = \left(\begin{array}{c|cccccc} & B_1 = N^+ \setminus \{g_0\} - N^- \setminus \{g_0^{-1}\} & g_0 - g_0^{-1} & B_2 & f^{(4)} & \mathcal{F}^2(f^{(4)}) \\ \hline N^+ \setminus \{g_0\} & I & 0 & I & \mathbf{1} & 0 \\ g_0 & 0 & 1 & -\mathbf{1} & 1 & 0 \\ N^- \setminus \{g_0^{-1}\} & -I & 0 & I & \mathbf{1} & 0 \\ g_0^{-1} & 0 & -1 & -\mathbf{1} & 1 & 0 \\ \mathcal{F}^2(f^{(4)}) & 0 & 0 & 0 & 0 & 1 \end{array} \right)$$

Setting $n = |G| - 1$, we have

$$(Q_D^B)^{-1} = \frac{1}{2n} \left(\begin{array}{c|ccc} & N^+ \setminus \{g_0\} & g_0 & N^- \setminus \{g_0^{-1}\} & g_0^{-1} & \mathcal{F}^2(f^{(4)}) \\ \hline B_1 & nI & 0 & -nI & 0 & 0 \\ g_0 - g_0^{-1} & 0 & n & 0 & -n & 0 \\ B_2 & nI - \mathbf{2} & -\mathbf{2} & nI - \mathbf{2} & -\mathbf{2} & 0 \\ f^{(4)} & \mathbf{2} & 2 & \mathbf{2} & 2 & 0 \\ \mathcal{F}^2(f^{(4)}) & 0 & 0 & 0 & 0 & 2n \end{array} \right)$$

Hence

$$[\mathcal{F}^2]_B = Q_D^B [\mathcal{F}^2]_D (Q_D^B)^{-1} = \frac{1}{n} \left(\begin{array}{c|cc} & N^+ & N^- & \mathcal{F}^2(f^{(4)}) \\ \hline N^+ & -\mathbf{1} & nI - \mathbf{1} & \mathbf{n} \\ N^- & nI - \mathbf{1} & -\mathbf{1} & \mathbf{n} \\ \mathcal{F}^2(f^{(4)}) & \mathbf{1} & \mathbf{1} & 0 \end{array} \right). \quad \square$$

Remark 2.14. Parts (1)-(3) of Conjecture A were chosen because they agree with 3^G for $G \in \{\mathbb{Z}/2, \mathbb{Z}/4, \mathbb{Z}/2 \times \mathbb{Z}/2\}$. There are two reasons +1 was chosen as the eigenvalue in part (4). First, if $S \neq \emptyset$ and $g, h \in N^+$ are distinct, then we have

$$(p_g + p_{g^{-1}}) - (p_h + p_{h^{-1}}) = 2p_{s_0} - (p_h + p_{h^{-1}}) - (2p_{s_0} - (p_g + p_{g^{-1}})),$$


which has eigenvalue 1 by part (3). Second, if $S = \emptyset$, switching the eigenvalue to -1 gives the formula

$$\mathcal{F}^2(p_g) = \frac{1}{n} \mathcal{F}^2(f^{(4)}) + \frac{n-1}{n} p_g - \frac{1}{n} \sum_{h \neq g} p_h,$$

which is not the same formula as when $S \neq \emptyset$.

3 The reduced subfactor at $\rho = f^{(2)}$

The even half \mathcal{E}_\bullet of \mathcal{P}_\bullet is a factor planar algebra [BHP12] with principal graph



We denote $\rho = f^{(2)}$ by a red strand, and we write a trivalent vertex for the intertwiner $f^{(2)} \otimes f^{(2)} \rightarrow f^{(2)}$ given by

$$\left[\begin{array}{c} \text{red strand} \\ \text{grey strand} \end{array} \right] = \left(\frac{[2]}{[3] - 1} \right)^{1/2} \left[\begin{array}{c} \text{grey strand} \\ \text{red strand} \end{array} \right] \quad (\text{V})$$

where we just write 2 for $f^{(2)}$.

Remark 3.1. It is a straightforward calculation that $|G| = \frac{[3]^2 - 1}{[3]}$.

Proposition 3.2. We have the following skein relations:

$$\begin{aligned}
 & \boxed{\text{circle with dot}} = \boxed{\text{two vertical lines}} \\
 & \boxed{\text{circle}} = \boxed{\text{circle with horizontal line}} = [3] \\
 & \boxed{\text{circle with dot}} = 0 \\
 & \boxed{\text{cup}} = \boxed{\text{cap}}^* = \boxed{\text{Y-junction}} \\
 & \boxed{\text{two vertical lines}} - \boxed{\text{cup and cap}} = ([3] - 1) \left(\boxed{\text{Y-junction}} - \boxed{\text{H-junction}} \right) \tag{I=H} \\
 & \boxed{\text{two vertical lines}} = \frac{1}{[3]} \boxed{\text{cup and cap}} + \boxed{\text{Y-junction}} + \sum_{g \neq 1} \boxed{p_g} \tag{E}
 \end{aligned}$$

Proof. The proof is similar to [IMP13, Proposition 3.1]. We prove Relations (I=H) and (E). To prove Relation (I=H), note $\dim(\text{Hom}(\rho^{\otimes 2}, \rho^{\otimes 2})) = 3$, so there is some linear relation amongst the four diagrams which appear in the relation. By rotational symmetry, we must have a relation of the form

$$\boxed{\text{two vertical lines}} \pm \boxed{\text{cup and cap}} = \lambda \left(\boxed{\text{Y-junction}} \pm \boxed{\text{H-junction}} \right).$$

We can determine λ by capping off the right hand side of all the diagrams. Finally, to determine that the sign is a minus sign, we note that the Temperley-Lieb diagram



has the same non-zero coefficient in both ρ -diagrams on the right when we expand the ρ -vertex and ρ -strands, but it does not appear on the left.

To prove Equation (E), we note that since $[2] > 2$, $f^{(2)} \otimes f^{(2)} \cong f^{(0)} \oplus f^{(2)} \oplus f^{(4)}$. Since $\sum_{g \neq 1} p_g = f^{(4)}$, we are finished. \square


Corollary 3.3. Since $|G| - 1 = \frac{[3]^2 - [3] - 1}{[3]}$, similar to Proposition 2.13, we have

$$\boxed{\text{H-junction}} = \frac{1}{|G| - 1} \left(\mathcal{F}^2(f^{(4)}) - f^{(4)} \right) + \boxed{\text{Y-junction}}.$$

Definition 3.4. We define $p_1 := \boxed{\text{Y-junction}} - \boxed{\text{H-junction}}$.

Definition 3.5. Let \mathcal{S}_\bullet be the reduced subfactor planar algebra of \mathcal{P}_\bullet at $\rho = f^{(2)}$. Since ρ is symmetrically self-dual in the sense of [MP15, MP], we may lift the shading on \mathcal{S}_\bullet to get a factor planar algebra \mathcal{R}_\bullet . We use the convention that $\mathcal{R}_n = \text{Hom}(\rho^{\otimes n}, \rho^{\otimes n})$, which is usually denoted \mathcal{R}_{2n} in [BHP12].

By Relation (E), the unitary fusion category associated to \mathcal{R}_\bullet is $\frac{1}{2}\mathcal{P}_+ \boxtimes \text{Vec}(\mathbb{Z}/2\mathbb{Z})$, and the principal graph Λ of \mathcal{R}_\bullet is given by




The principal graphs of \mathcal{S}_\bullet are (Λ, Λ) , and $\frac{1}{2}\mathcal{S}_+ = \frac{1}{2}\mathcal{P}_+$.

Remarks 3.6.

- (1) We may also obtain \mathcal{S}_\bullet by imposing a shading on \mathcal{E}_\bullet .
- (2) We may naturally identify $\mathcal{R}_n \cong \mathcal{S}_{n,\pm}$ as a subspace of $\mathcal{P}_{2n,+}$.

Proposition 3.7. *Suppose that $A \in \mathcal{R}_2$ is rectangularly uncappable in $\mathcal{P}_{4,+}$, i.e., capping A on the top or bottom is zero when we write A with 4 strands up and 4 strands down. Then we may identify the two click rotation $\mathcal{F}_{\mathcal{P}_\bullet}^2(A) \in \mathcal{P}_{4,+}$ with the 1-click rotation $\mathcal{F}_{\mathcal{R}_\bullet}(A) \in \mathcal{R}_2$.*



Consider the orthogonal complement of $\mathcal{TL}_{4,+} \subset \mathcal{P}_{4,+}$, which is spanned by $|G| - 2$ low weight vectors $\{A_j | j = 1, \dots, |G| - 2\} \subset \text{span}\{p_g | g \neq 1\}$, which are also eigenvectors for the 2-click rotation \mathcal{F}^2 on \mathcal{P}_\bullet corresponding to rotational eigenvectors ω_{A_j} . By Proposition 3.7, we get the following corollary:

Corollary 3.8. *Each A_j is also a low-weight rotational eigenvector for the 1-click rotation in \mathcal{R}_\bullet .*

Since $\sum_{g \neq 1} p_g = f^{(4)}$, we know that another low weight rotational eigenvector in $\mathcal{R}_{2,+}$ comes from $\mathcal{TL}_{4,+} \subset \mathcal{P}_{4,+}$:

$$B = (|G| - 1)p_1 - f^{(4)} = (|G| - 1)p_1 - \sum_{g \neq 1} p_g,$$

where the rotational eigenvalue $\omega_B = 1$ by Corollary 3.3. Note that B is orthogonal to each A_j .

We now compute the 1-click rotation for all the non-trivial minimal projections in \mathcal{R}_2 in terms of the p_g for $g \in G$ and

$$e_1 = \frac{1}{[3]} \left[\text{diagram} \right] \text{ and } f^{(2)} = \left[\text{diagram} \right] - e_1 = \left[\text{diagram} \right] + \sum_{g \neq 1} \left[\text{diagram} \right].$$

The diagrams in the equation are: a box with two crossings, a box with two vertical strands, a box with two crossings, and a box with two vertical strands and a red double line.

Note that by Relations (I=H) and (E),

$$\begin{aligned} \mathcal{F}^2(f^{(4)}) &= \frac{1}{[3]([3] - 1)} f^{(4)} + \frac{[3]^2 - [3] - 1}{[3]} e_1 - \frac{[3]^2 - [3] - 1}{[3]([3] - 1)} p_1 \\ &= \frac{1}{[3]([3] - 1)} f^{(4)} + (|G| - 1)e_1 - \frac{|G| - 1}{[3] - 1} p_1. \end{aligned}$$

Corollary 3.9. *Conjecture A holds if and only if for each $g \in G$,*

$$\begin{aligned}
 \mathcal{F}_{\mathcal{R}_\bullet}(p_g) &= \frac{1}{|G|-1}(\mathcal{F}^2(f^{(4)}) - f^{(4)}) + p_{g^{-1}} \\
 &= p_{g^{-1}} - \frac{1}{[3]-1}f^{(4)} + e_1 - \frac{1}{[3]-1}p_1 \\
 &= p_{g^{-1}} + e_1 - \frac{1}{[3]-1} \sum_{g \in G} p_g \\
 &= p_{g^{-1}} + e_1 - \frac{1}{[3]-1}f^{(2)} \\
 &= p_{g^{-1}} - \frac{1}{[3]-1} \left(\begin{array}{|c|} \hline \text{---} \\ \hline \end{array} - \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \right).
 \end{aligned}$$

4 An ‘almost’ G -action on \mathcal{R}_\bullet .

We now construct an ‘almost’ G -action on \mathcal{R}_\bullet . This corresponds to an action of G on $\frac{1}{2}\mathcal{P}_+$. We define for each $g \in G$ a map Φ_g on the unshaded factor planar algebra \mathcal{R}_\bullet .

Definition 4.1. Recall that the group G can be seen in $\frac{1}{2}\mathcal{P}_+$ inside $\mathcal{P}_{6,+}$ as the minimal projections at depth 6 of Γ_+ , together with the image of the trivial object in $\mathcal{P}_{6,+}$ given by

$$e = \frac{1}{[4]} \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \text{ where } \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} = f^{(3)} \in \mathcal{P}_{3,+}.$$

Here, we switch the convention of the unit of the group to e instead of 1 to not confuse the empty diagram 1 with the projection $e \in \mathcal{P}_{6,+}$. (While the empty diagram is the identity of $\mathcal{P}_{0,+}$, e is not the identity of $\mathcal{P}_{6,+}$!)

The group multiplication is given by a multiple of the coproduct.

Lemma 4.2. *In $\mathcal{P}_{6,+}$, the coproduct $g * h = \begin{array}{|c|} \hline g \\ \hline \end{array} \begin{array}{|c|} \hline h \\ \hline \end{array} = [4]^{-1} \begin{array}{|c|} \hline gh \\ \hline \end{array}.$*

Proof. We know $g \otimes h \cong gh$ and $g * h$ is self adjoint, so $g * h = \lambda gh$ for some $\lambda \in \mathbb{R}$. Taking traces shows that $\lambda = [4]^{-1}$. □

Recall that $\rho g \cong g^{-1}\rho$ for all $g \in G$. This means there is exactly one non-zero morphism up to scaling between the two.

Definition 4.3. For $g \in G$, we define the following element of $\mathcal{P}_{6,+}$:

$$V_g = [3] \begin{array}{|c|} \hline g \\ \hline \end{array} \begin{array}{|c|} \hline g \\ \hline \end{array} = [3] \begin{array}{|c|} \hline g \\ \hline \end{array} \begin{array}{|c|} \hline g \\ \hline \end{array} = \begin{array}{|c|} \hline g \\ \hline \end{array} \begin{array}{|c|} \hline g \\ \hline \end{array}.$$

We may think of V_g as a crossing as above, where we use an oriented strand labelled by g to denote three ρ -strands cabled by g , and the direction gives the location of the \star . Here, we use the convention that V_e is given by


$$V_e = \frac{1}{[4]} \begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} \cup \\ \cap \end{array}.$$

Note that we also have

$$V_g^* = [3] \begin{array}{c} \star \\ \boxed{g} \\ \star \end{array} \begin{array}{c} \star \\ \boxed{g} \\ \star \end{array} = [3] \begin{array}{c} \star \\ \boxed{g} \\ \star \end{array} \begin{array}{c} \star \\ \boxed{g} \\ \star \end{array} = \star \begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} \cup \\ \cap \end{array} \star.$$

Using V_g and V_g^* , we define the map Φ_g on $x \in \mathcal{R}_n$ by encircling x by a strand whose orientation reverses as it crosses the ρ -strands connected to x . The orientation is clockwise in the distinguished region of x . We replace crossings with either V_g or V_g^* depending on the crossings. This means that if we travel on the g -strand from an unshaded region to a shaded region, we replace the crossing with V_g , and if we cross from shaded to unshaded we replace the crossing with V_g^* .

Remark 4.4. It is easy to see that for $x \in \mathcal{R}_n$ and $g \in G$, $\Phi_g(x) \in \mathcal{R}_n$. When $g = e$, Φ_e is the identity. When $g \neq e$, $V_g, V_g^* \in \mathcal{R}_4$, so $\Phi_g(x) \in \mathcal{R}_n$.



Example 4.5. When $x \in \mathcal{R}_2$, we have $\Phi_g(x) =$


Lemma 4.6. There is a constant $\theta_g \in U(1)$ such that $\mathcal{F}(V_g^*) = \theta_g V_{g^{-1}}$ and $\mathcal{F}^{-1}(V_g) = \theta_g^{-1} V_{g^{-1}}^*$. Moreover, $\theta_g = \theta_{g^{-1}}$ for all $g \in G$, and $\theta_e = 1$.

Proof. There is exactly one map up to a scalar from $g^{-1}\rho$ to ρg , so there is a constant $\theta_g \neq 0$ such that $\mathcal{F}(V_g^*) = \theta_g V_{g^{-1}}$. Since the norm squared of $\mathcal{F}(V_g^*)$ equals the norm squared of $V_{g^{-1}}$, $\theta_g \in U(1)$. Now taking adjoints, we have $\mathcal{F}^{-1}(V_g) = \theta_g^{-1} V_{g^{-1}}^*$.

We now apply \mathcal{F} to the equation $\mathcal{F}^{-1}(V_g) = \theta_g^{-1} V_{g^{-1}}^*$ to get the equation $\mathcal{F}(V_{g^{-1}}^*) = \theta_g V_g$. This means $\theta_g = \theta_{g^{-1}}$ by the definition of $\theta_{g^{-1}}$.

Finally, a simple diagrammatic calculation shows $\theta_e = 1$. □

Corollary 4.7. The map Φ is also given as follows. First, encircle x by a strand whose orientation reverses as it crosses the ρ -strands connected to x . The orientation is clockwise in the distinguished region of x . We replace crossings with either $V_{g^{-1}}$ or $V_{g^{-1}}^*$ depending on the crossings.



Corollary 4.8. We have $\Phi_g \circ \mathcal{F}_{\mathcal{R}_\bullet} = \mathcal{F}_{\mathcal{R}_\bullet} \circ \Phi_{g^{-1}}$.


Remark 4.9. Since each $g \in G$ has dimension 1, we have the usual skein relations for g -strands:

$$\begin{array}{|c|} \hline \text{---} \\ \hline \end{array} = \begin{array}{|c|} \hline \downarrow \\ \hline \end{array} \text{ and } \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} = \begin{array}{|c|} \hline \downarrow \\ \hline \end{array}.$$

Proposition 4.10. The g -strand and the ρ -strand together with $V_g, V_g^*, V_{g^{-1}}, V_{g^{-1}}^*$, satisfy the Reidemeister II relations

$$(1) \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} = \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \text{ and } (2) \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} = \begin{array}{|c|} \hline \text{---} \\ \hline \end{array}.$$

Proof. The case $g = e$ is trivial. When $g \neq e$, we prove (1), and (2) follows by replacing g with g^{-1} . The second equality follows from the fact that each of $V_g, V_g^*, V_{g^{-1}}, V_{g^{-1}}^*$ is fixed under $\mathcal{F}_{\mathcal{R}_\bullet}^4$. To prove the first equality in (1), we see



using the skein relation from Remark 4.9 for g -cabled strands. □

Proposition 4.11. The map Φ_g is compatible with the graded multiplication operator given for $x \in \mathcal{R}_m$ and $y \in \mathcal{R}_n$ by

$$x \wedge y = \begin{array}{|c|} \hline m \\ \hline x \\ \hline \end{array} \begin{array}{|c|} \hline n \\ \hline y \\ \hline \end{array}.$$

Proof. Just use the skein relation in Remark 4.9:

$$\Phi_g(x) \wedge \Phi_g(y) = \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} = \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} = \Phi_g(x \wedge y). \quad \square$$

Corollary 4.12. The ‘almost’ action of Φ_g on \mathcal{R}_\bullet induces the action of g^2 on $\frac{1}{2}\mathcal{P}_+$.

Proof. For $h, g \in G$, we will show that $\Phi_g(p_h \in \mathcal{S}_{2,+}) = p_{g^2h}$ and $\Phi_g(h \in \mathcal{S}_{3,+}) \cong g^2h$. As both proofs are similar, we will only show $\Phi_g(p_h) = p_{g^2h}$.

First, by Propositions 4.10 and 4.11 together with the fact that h is an orthogonal projection, we see $\Phi_g(p_h)^2 = \Phi_g(p_h) = \Phi_g(p_h)^*$, i.e., $\Phi_g(p_h)$ is an orthogonal projection. Next, taking the trace, we see $\Phi_g(p_h) \neq 0$ by sphericity and again using the Reidemeister II relation from Proposition 4.10:

Finally, it's obvious that $\Phi_g(p_h) \cong g \otimes (h\rho) \otimes g^{-1} \cong g^2 h\rho$, so $\Phi_g(p_h) = p_{g^2h}$ since \mathcal{R}_2 is abelian. □

Proposition 4.13. For every $g \in G$, we have

$$\mathcal{F}_{\mathcal{R}_\bullet}(p_{g^2}) = p_{g^{-2}} - \frac{1}{[3] - 1} \left(\begin{array}{|c|} \hline \text{[Diagram: square with two arcs]} \\ \hline \end{array} - \begin{array}{|c|} \hline \text{[Diagram: square with two vertical lines]} \\ \hline \end{array} \right).$$

(Compare the above equation with Corollary 3.9.)

Proof. By Relation I=H, we know

$$\mathcal{F}_{\mathcal{R}_\bullet}(p_1) = p_1 - \frac{1}{[3] - 1} \left(\begin{array}{|c|} \hline \text{[Diagram: square with two arcs]} \\ \hline \end{array} - \begin{array}{|c|} \hline \text{[Diagram: square with two vertical lines]} \\ \hline \end{array} \right).$$

Apply Φ_g to the equation to see

$$\begin{aligned} \mathcal{F}_{\mathcal{R}_\bullet}(p_{g^2}) &= \mathcal{F}_{\mathcal{R}_\bullet}(\Phi_g(p_1)) \\ &= \Phi_{g^{-1}}(\mathcal{F}_{\mathcal{R}_\bullet}(p_1)) \\ &= \Phi_{g^{-1}} \left(p_1 - \frac{1}{[3] - 1} \left(\begin{array}{|c|} \hline \text{[Diagram: square with two arcs]} \\ \hline \end{array} - \begin{array}{|c|} \hline \text{[Diagram: square with two vertical lines]} \\ \hline \end{array} \right) \right) \\ &= \Phi_{g^{-1}}(p_1) - \frac{1}{[3] - 1} \left(\begin{array}{|c|} \hline \text{[Diagram: square with two arcs]} \\ \hline \end{array} - \begin{array}{|c|} \hline \text{[Diagram: square with two vertical lines]} \\ \hline \end{array} \right) \\ &= p_{g^{-2}} - \frac{1}{[3] - 1} \left(\begin{array}{|c|} \hline \text{[Diagram: square with two arcs]} \\ \hline \end{array} - \begin{array}{|c|} \hline \text{[Diagram: square with two vertical lines]} \\ \hline \end{array} \right), \end{aligned}$$

where we used $\Phi_{g^{-1}} \circ \mathcal{F}_{\mathcal{R}_\bullet} = \mathcal{F}_{\mathcal{R}_\bullet} \circ \Phi_g$ by Corollary 4.8. □

We can now prove our main theorem, which says that Conjecture A is true for $|G|$ odd.

Proof of Theorem B. Note that the action induced by Φ of G on $G\rho = \{h\rho | h \in G\}$ is freely transitive exactly when $|G|$ is odd. This is because the action is given by $\Phi_g(h\rho) = g^2 h\rho$ and $\Phi_g(h) = g^2 h$ by Corollary 4.12, and the map $g \mapsto g^2$ is an automorphism of G when G is odd.

Thus for every $g \in G$, the equation in Corollary 3.9 holds by Proposition 4.13, which concludes the proof. □

4.1 Lifting involutions to the center

Definition 4.14. Let \mathcal{Q}_\bullet be the planar subalgebra of \mathcal{R}_\bullet generated by

$$\left\{ \begin{array}{|c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \left| g \neq 1 \right. \right\} \text{ and } p_1 := \begin{array}{|c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array},$$

under the ρ -strand planar operad, whose tangles do not contain trivalent vertices.

We expect the following conjecture to be true, but it seems to be highly non-trivial at this time. We will prove it for the case $|G|$ is odd in Theorem 5.19.


Conjecture 4.15. $\mathcal{Q}_\bullet = \mathcal{R}_\bullet$.

We now show that each involution in G , i.e., a $g \in G$ with $g^2 = 1$, lifts to the center of the projection category of \mathcal{Q}_\bullet , i.e., $\mathcal{Z}(\text{Pro}(\mathcal{Q}_\bullet))$ [BHP12]. By Corollary 4.12, each Φ_g restricts to an automorphism of \mathcal{Q}_\bullet .

Lemma 4.16. *If $g^2 = 1$, then $\Phi_g = \text{id}_{\mathcal{Q}_\bullet}$.*

Proof. In the proof of Corollary 4.12, we showed that $\Phi_g(p_h) = p_{g^2h}$ for all $h \in H$. Since $g^2 = 1$, Φ_g fixes every p_h , which generate \mathcal{Q}_\bullet as a planar algebra. □


Proposition 4.17. *For all $\varphi \in \mathcal{Q}_n$, we have*



where the \bullet is either blank or $*$ depending on the parity of n .


Proof. Take the norm squared of the difference and use Lemma 4.16. □

Corollary 4.18. *Suppose $g \in G$ is an involution. The map $e_g : X \rightarrow \text{Hom}(g \otimes X, X \otimes g)$ for simple X by $e_g(\rho) = V_g$ and*



naturally extends to a half-braiding. Hence (g, e_g) defines an element in the center $\mathcal{Z}(\text{Pro}(\mathcal{Q}_\bullet))$.

Corollary 4.19. *Suppose $g \in G$ is an involution. The twist factor of (g, e_g) is given by θ_g .*



5 The G -action on \mathcal{S}_\bullet and bases for $\mathcal{S}_{3,+}$

We saw each Φ_g almost gives an automorphism of the unshaded factor planar algebra \mathcal{R}_\bullet , except for the problem with the 1-click rotation from Corollary 4.8. However, Φ_g is compatible with the 2-click rotation, which suggests that the Φ_g 's can be used to construct automorphisms of the shaded subfactor planar algebra \mathcal{S}_\bullet . There is a slight technicality here – when we identify $\mathcal{S}_{n,\pm}$ with \mathcal{R}_n , the map Φ_g goes from $\mathcal{S}_{n,\pm} \rightarrow \mathcal{S}_{n,\mp}$. Hence to get a planar algebra map, we need to also use the symmetric self-duality $\Delta_\pm : \mathcal{S}_\pm \rightarrow \mathcal{S}_\mp$ which reverses the shading.

Definition 5.1. For $g \in G$, define Ψ_g on $\mathcal{S}_{n,\pm} = \mathcal{R}_n$ by $\Psi_g = \Delta_\mp \circ \Phi_{g^\pm 1}$.

Example 5.2. When $x \in \mathcal{S}_{2,+}$ and $y \in \mathcal{S}_{3,-}$, we have

$$\Psi_g(x) = \Delta_- \left(\begin{array}{c} \text{Diagram 1: } x \text{ in } \mathcal{S}_{2,+} \end{array} \right) = \Delta_- \left(\begin{array}{c} \text{Diagram 2: } x \text{ in } \mathcal{S}_{2,-} \end{array} \right) \text{ and}$$

$$\Psi_g(y) = \Delta_+ \left(\begin{array}{c} \text{Diagram 3: } y \text{ in } \mathcal{S}_{3,-} \end{array} \right) = \Delta_+ \left(\begin{array}{c} \text{Diagram 4: } y \text{ in } \mathcal{S}_{3,+} \end{array} \right).$$

Lemma 5.3. On \mathcal{S}_- , $\Psi_g = \mathcal{F}_{\mathcal{S}_\bullet}^{-1} \circ \Psi_g \circ \mathcal{F}_{\mathcal{S}_\bullet}$.

Proof. Identifying $\mathcal{S}_{n,\pm} = \mathcal{R}_n$, we have $\mathcal{F}_{\mathcal{S}_\bullet} = \mathcal{F}_{\mathcal{R}_\bullet}$. Since Δ is a symmetric self-duality, $\Delta_+ \circ \mathcal{F}_{\mathcal{S}_\bullet} = \mathcal{F}_{\mathcal{S}_\bullet} \circ \Delta_-$. By Corollary 4.8, $\mathcal{F}_{\mathcal{R}_\bullet} \circ \Phi_{g^{-1}} = \Phi_g \circ \mathcal{F}_{\mathcal{R}_\bullet}$. Combining these, we have

$$\Psi_g = \Delta_+ \circ \Phi_{g^{-1}} = (\Delta_+ \circ \mathcal{F}_{\mathcal{R}_\bullet}^{-1}) \circ (\mathcal{F}_{\mathcal{R}_\bullet} \circ \Phi_{g^{-1}}) = \mathcal{F}_{\mathcal{R}_\bullet}^{-1} \circ (\Delta_- \circ \Phi_g) \circ \mathcal{F}_{\mathcal{R}_\bullet} = \mathcal{F}_{\mathcal{S}_\bullet}^{-1} \circ \Psi_g \circ \mathcal{F}_{\mathcal{S}_\bullet}. \quad \square$$



Proposition 5.4. The map Ψ_g is a shaded planar algebra automorphism.

Proof. We need to show that Ψ_g commutes with a set of generating tangles for the planar operad. We can use the graded multiplication operator and the annular tangles, which are clearly generated by adding cups and caps and the 1-click rotation operator.

The fact that Ψ_g is compatible with the 1-click rotation is exactly Lemma 5.3. We see that Ψ_g is compatible with cups and caps by the Reidemeister II relation from Proposition 4.10. Finally, Ψ_g is compatible with the graded multiplication operator by Proposition 4.11. \square

5.1 The G -action on \mathcal{S}_\bullet

We prove a few lemmas to calculate constants, after which we will see Ψ gives an action of G on \mathcal{S}_\bullet .

Corollary 5.5.  and .

Proof. The first equation follows by taking the norm squared of the difference and applying Lemma 4.2. The second equation then follows immediately. \square


Definition 5.6. For $h, g \in G$, let the (g, h, hg) -trivalent vertex be given by

$$Y_{g,h} = \begin{array}{c} gh \\ \diagup \quad \diagdown \\ g \quad h \end{array} = [4]^{1/2} \begin{array}{c} gh \\ \diagup \quad \diagdown \\ g \quad h \end{array},$$


so that $Y_{g,h}^* Y_{g,h} = gh$ by Corollary 5.5. Since $Y_{g,h} Y_{g,h}^*$ also has trace 1, we immediately have that

$$g \uparrow h \downarrow = \begin{array}{c} g \quad h \\ \diagdown \quad \diagup \\ gh \end{array} = [4] \begin{array}{c} gh \\ \diagup \quad \diagdown \\ g \quad h \end{array} = \begin{array}{c} g \quad h \\ \diagdown \quad \diagup \\ gh \end{array}. \tag{2}$$

Remark 5.7. Note that the (g, h, gh) -trivalent vertex $Y_{g,h}$ is not in \mathcal{R}_\bullet . At this point, we do not know whether $Y_{g,h} Y_{g,h}^*$ is in \mathcal{R}_\bullet .

Lemma 5.8. The G -trivalent vertices are associative, i.e., .

Proof. Using Corollary 5.5, we have



Remark 5.9. Suppose we did not assume that \mathcal{R}_\bullet was the reduced subfactor planar algebra of \mathcal{P}_\bullet at $\rho = f^{(2)}$, and instead we started with a factor planar algebra with principal graph Λ from Definition

3.5. In this case, we might have that the G -trivalent vertices are not associative. Rather, there may be a non-trivial 3-cocycle giving a non-trivial associator for G .

In fact, there are such examples for $\mathbb{Z}/3\mathbb{Z}$ giving ‘twisted’ Haagerup categories due to Ostrik [MPS15, Proposition 7.7, and the following paragraph].

Recall from Lemma 4.6 that there is a distinguished 1-cochain $\theta \in C^1(G, U(1))$.

Corollary 5.10. *For all $g, h \in G$, there is a scalar $\mu_{g,h} \in U(1)$ such that*

Moreover, $\mu \in Z^2(G, U(1))$, and $\mu_{g,h}\mu_{g^{-1},h^{-1}} = [(d\theta)(g, h)]^{-1}$.

Proof. Take the norm squared of each diagram, unzip the trivalent vertices, and use the Reidemeister II relation from Proposition 4.10 to get that both closed diagrams equal [3]. Since there is only one map up to scaling from $gh \otimes \rho$ to $\rho \otimes (gh)^{-1}$, both sides must be equal up to a phase, denoted $\mu_{g,h}$.

A straightforward calculation again by unzipping and using the Reidemeister II relation from Proposition 4.10 shows that for $g, h, k \in G$, $\mu_{g,hk}\mu_{h,k} = \mu_{gh,k}\mu_{g,h}$, i.e., $\mu \in Z^2(G, U(1))$.

For the final claim, we first look at

This must be equal to

This means that we must have $\mu_{g,h}\mu_{g^{-1},h^{-1}} = (\theta_g\theta_{g^{-1}}\theta_h)^{-1} = [(d\theta)(g, h)]^{-1}$. □

Remark 5.11. The significance of the final formula in Corollary 5.10 is that there is a strong relation between the structure constants of \mathcal{S}_\bullet .

Corollary 5.12. *On \mathcal{R}_n , we have $\Phi_g \circ \Phi_h = \Phi_{gh}$, and similarly for the Ψ_g ’s on $\mathcal{S}_{n,\pm}$.*

Proof. Given $x \in \mathcal{S}_{n,+}$, start with the diagram for $\Phi_g \circ \Phi_h(x)$, use Relation (2), and apply Corollary 5.10. We will get alternating contributions of $\mu_{g,h}$ and $\overline{\mu_{g,h}} = \mu_{g,h}^{-1}$, which cancel, leaving us with the diagram for $\Phi_{gh}(x)$. The proof for $y \in \mathcal{S}_{n,-}$ is similar. □

5.2 Bases for $\mathcal{S}_{3,+}$

We now define some distinguished elements for $\mathcal{S}_{3,+}$. When $|G|$ is odd, we show these elements form a basis for $\mathcal{S}_{3,+}$. We then prove Theorem D and Conjecture 4.15 in the case $|G|$ is odd.

We use the notation $p_\emptyset = e_1 = \frac{1}{[3]} \begin{array}{|c|} \hline \square \\ \hline \end{array}$.

Definition 5.13. For $i, j \in G \cup \{\emptyset\}$ and $g, h, k \in G$, define

$$\alpha_{i,j} = \frac{1}{\text{Tr}(p_i)^{1/2} \text{Tr}(p_j)^{1/2}} \begin{array}{|c|} \hline p_j \\ \hline p_i \\ \hline \end{array} \quad \text{and} \quad \beta_{h,k,\ell} = \begin{array}{|c|} \hline \star(p_\ell) \\ \hline \star(p_h) \\ \hline \end{array} \begin{array}{|c|} \hline \star(p_k) \\ \hline \end{array}.$$

The following facts are straightforward.

Facts 5.14.

- (1) The elements $\{\alpha_{i,j}\}$ form a system of matrix units for the copy of $M_{|G|+1}(\mathbb{C})$ corresponding to $\mathcal{I}_{3,+} = \mathcal{S}_{2,+} e_2 \mathcal{S}_{2,+}$.
- (2) The inner product $\langle \beta_{g,h,k}, \beta_{g',h',k'} \rangle$ is zero unless $g = g', h = h',$ and $k = k'$.
- (3) For all $g \in G$, we have $\Psi_g(\alpha_{i,j}) = \alpha_{g^2 i, g^2 j}$, where we define $g^2 \emptyset = \emptyset$.
- (4) For all $g \in G$, we have $\Psi_g(\beta_{h,k,\ell}) = \beta_{g^2 h, g^2 k, g^2 \ell}$.

Lemma 5.15. If $P_{\mathcal{I}_{3,+}}$ is the orthogonal projection onto $\mathcal{I}_{3,+}$ in $\mathcal{S}_{3,+}$, then $P_{\mathcal{I}_{3,+}}(\beta_{h,k,\ell}) = c_{h,k,\ell} \alpha_{h,\ell}$ where

$$c_{h,k,\ell} = \frac{1}{[3]} \begin{array}{|c|} \hline \star(p_\ell) \\ \hline \star(p_h) \\ \hline \end{array} \begin{array}{|c|} \hline \star(p_k) \\ \hline \end{array}.$$

Proof. Using (1) from Facts 5.14, we see that the projection of $\beta_{h,k,\ell}$ onto $\mathcal{I}_{3,+}$ is given by

$$P_{\mathcal{I}_{3,+}}(\beta_{h,k,\ell}) = \sum_{i,j} \frac{\langle \beta_{h,k,\ell}, \alpha_{i,j} \rangle}{\|\alpha_{i,j}\|_2^2} \alpha_{i,j} = \sum_{i,j} \frac{\text{Tr}(\beta_{h,k,\ell} \alpha_{i,j}^*)}{\text{Tr}(\alpha_{i,j} \alpha_{i,j}^*)} \alpha_{i,j} = \frac{\text{Tr}(\beta_{h,k,\ell} \alpha_{h,\ell}^*)}{\text{Tr}(\alpha_{h,\ell} \alpha_{h,\ell}^*)} \alpha_{h,\ell} = c_{h,k,\ell} \alpha_{h,\ell}. \quad \square$$


Lemma 5.16. If k has a square root in G , then

$$c_{h,k,\ell} = \frac{1}{[3]} \begin{array}{|c|} \hline \star(p_\ell) \\ \hline \star(p_h) \\ \hline \end{array} \begin{array}{|c|} \hline \star(p_k) \\ \hline \end{array} = \frac{1}{[3]} \begin{array}{|c|} \hline \star(p_{k^{-1}\ell}) \\ \hline \star(p_{k^{-1}h}) \\ \hline \end{array} \begin{array}{|c|} \hline \star(p_1) \\ \hline \end{array} = \begin{cases} \frac{[3]}{([3]-1)^2} & \text{if } h, k, \ell \text{ are distinct} \\ 1 & \text{if any two are equal} \\ \left(\frac{[3]-2}{[2]}\right)^2 & \text{if } h = k = \ell. \end{cases}$$

By symmetry, a similar statement holds if h or ℓ has a square root in G .

Moreover, if any of h, k, ℓ have a square root in G , then $\gamma_{h,k,\ell} := \beta_{h,k,\ell} - c_{h,k,\ell} \alpha_{h,\ell} \neq 0$.

Proof. Using (4) of Facts 5.14 and sphericity, we have

$$\langle \beta_{h,k,\ell}, \alpha_{h,\ell} \rangle = \|\beta_{h,k,\ell}\|_2^2 = \|\beta_{g^2(g^{-2}h), g^2, g^2(g^{-2}\ell)}\|_2^2 = \|\Psi_g(\beta_{k^{-1}h, 1, k^{-1}\ell})\|_2^2 =$$


When $h \neq k \neq \ell$, expanding p_1 using Equation (V) and simplifying, we get

$$\|\beta_{h,k,\ell}\|_2^2 = \left(\frac{[2]}{[3]-1} \right) \left(\frac{[3]^2}{[4]} - \delta_{h,\ell} \frac{[3]}{[2]} \right) = \begin{cases} \frac{[2][3]^2}{([3]-1)[4]} & \text{if } h \neq \ell \\ \frac{[2]^2[3]}{([3]-1)[2][4]} & \text{if } h = \ell \end{cases} = \begin{cases} \frac{[3]^2}{([3]-1)^2} & \text{if } h \neq \ell \\ \frac{[3]}{([3]-1)^2} & \text{if } h = \ell. \end{cases}$$

A similar calculation handles the cases $h = k \neq \ell$ and $h \neq k = \ell$.

Finally, for the case $h = k = \ell$, the following relation derived using Equation (V) is helpful:

$$\boxed{\text{triangle}} = \left([2] - \frac{3}{[2]} \right) \boxed{\text{triangle}}.$$

Again using Ψ_g , we see $c_{h,h,h}$ is a multiple of the inner product of two triangles:

$$c_{h,h,h} = \frac{1}{[3]} \langle \boxed{\text{triangle}}, \boxed{\text{triangle}} \rangle = \left([2] - \frac{3}{[2]} \right)^2 = \left(\frac{[3]-2}{[2]} \right)^2.$$

To prove $\gamma_{h,k,\ell} \neq 0$, a straightforward calculation shows $\|\alpha_{h,\ell}\|_2^2 = [3]^{-1}$, which implies

$$\|c_{h,k,\ell}\alpha_{h,\ell}\|_2^2 = \frac{|c_{h,k,\ell}|^2}{[3]} \neq \|\beta_{h,k,\ell}\|_2^2. \quad \square$$

Proposition 5.17. *When $|G|$ is odd, $\{\gamma_{h,k,\ell} = \beta_{h,k,\ell} - c_{h,\ell}\alpha_{h,\ell}\}$ is a basis of $\mathcal{S}_{3,+} \ominus \mathcal{I}_{3,+}$.*

Proof. By counting dimensions, it suffices to show linear independence. Suppose we have a linear combination

$$0 = \sum_{h,k,\ell \in G} \lambda_{h,k,\ell} \gamma_{h,k,\ell}.$$

Compress by a particular p_h and p_ℓ on the bottom and top respectively to get

$$0 = \sum_{k \in G} \lambda_{h,k,\ell} \gamma_{h,k,\ell}.$$

Attaching by p_g on the right hand side, we see for every $g \in G$,

$$0 = \sum_{k \in G} \lambda_{h,k,\ell} \boxed{\text{triangle}} \boxed{p_g} = \lambda_{h,g,\ell} \beta_{h,g,\ell} - \sum_{k \in G} \lambda_{h,k,\ell} c_{h,\ell} \beta_{h,g,\ell} = \left(\lambda_{h,g,\ell} - \sum_{k \in G} \lambda_{h,k,\ell} c_{h,\ell} \right) \beta_{h,g,\ell}.$$

Hence $\lambda_{h,g,\ell} = \sum_{k \in G} \lambda_{h,k,\ell} c_{h,\ell}$ is independent of $g \in G$. Denote this common value by $\lambda_{h,\ell}$. We now see that $(1 - |G|c_{h,\ell})\lambda_{h,\ell} = 0$, which implies $\lambda_{h,\ell} = 0$. (A straightforward calculation using Remark 3.1 and Lemma 5.16 shows $c_{h,\ell} \neq |G|^{-1}$.) \square

Corollary 5.18. *When $|G|$ is odd, $\mathcal{B}_{3,+} = \{\alpha_{i,j}\}_{i,j \in G \cup \{\emptyset\}} \cup \{\gamma_{h,k,\ell}\}_{h,k,\ell \in G}$ is a basis of $\mathcal{S}_{3,+}$.*

Recall that \mathcal{Q}_\bullet is the planar subalgebra of \mathcal{R}_\bullet generated by \mathcal{R}_2 . We now prove Conjecture 4.15 for the case when $|G|$ is odd.

Theorem 5.19. *When $|G|$ is odd, $\mathcal{Q}_\bullet = \mathcal{R}_\bullet$.*

Proof. First, identifying $\mathcal{S}_{3,+}$ with \mathcal{R}_3 , we note that every element in $\mathcal{B}_{3,+}$ is in \mathcal{Q}_3 . Using Wenzl’s generalized relation [Liu13, Section 2.3], we see that there is a basis of \mathcal{R}_4 of elements in \mathcal{Q}_4 , since every element in the basic construction ideal $\mathcal{I}_4 = \mathcal{R}_3 e_3 \mathcal{R}_3$ is in \mathcal{Q}_4 . Since \mathcal{R}_\bullet has depth 4, all further box spaces are equal to the corresponding basic construction ideals, and thus $\mathcal{Q}_\bullet = \mathcal{R}_\bullet$.

We give an alternate argument. Since $\mathcal{B}_{3,+} \subset \mathcal{Q}_3$, $\mathcal{Q}_3 = \mathcal{R}_3$, so the principal graphs agree to depth 3. By counting dimensions, there is only one way for this graph to terminate, so the principal graphs of \mathcal{Q}_\bullet and \mathcal{R}_\bullet are equal, and $\mathcal{Q}_\bullet = \mathcal{R}_\bullet$. \square

Theorem (Theorem D). *For $|G|$ odd, \mathcal{S}_\bullet is a Yang-Baxter planar algebra with $|G| - 1$ generators.*

Proof. By Theorem 5.19, \mathcal{S}_\bullet is generated by 2-boxes. For $i, j \in G \cup \{\emptyset\}$ and $h, k, \ell \in G$, we may similarly define elements by considering

$$\eta_{i,j} = \frac{1}{\text{Tr}(p_i)^{1/2} \text{Tr}(p_j)^{1/2}} \text{ and } \xi_{h,k,\ell} = \text{diagram}$$

Similar calculations show $\mathcal{B}'_{3,+} = \{\eta_{i,j}\}_{i,j \in G \cup \{\emptyset\}} \cup \{\zeta_{h,k,\ell} = \xi_{h,k,\ell} - c_{h,\ell} \eta_{h,\ell}\}_{h,k,\ell \in G}$ is a basis for $\mathcal{S}_{3,+} \ominus \mathcal{I}_{3,+}$.

Now for all $h, k, \ell \in G$, $\xi_{h,k,\ell} \in \text{span}(\mathcal{B}_{3,+})$ and $\beta_{h,k,\ell} \in \text{span}(\mathcal{B}'_{3,+})$, which gives the necessary relations to show \mathcal{S}_\bullet is a Yang-Baxter planar algebra. \square

Remark 5.20. At this point, it seems highly non-trivial to compute the structure constants $\langle \beta_{h,k,\ell}, \xi_{h',k',\ell'} \rangle$.

References

- [AH99] Marta Asaeda and Uffe Haagerup, *Exotic subfactors of finite depth with Jones indices $(5 + \sqrt{13})/2$ and $(5 + \sqrt{17})/2$* , Comm. Math. Phys. **202** (1999), no. 1, 1–63, [arXiv:math.OA/9803044](#) [MR1686551](#) [DOI: 10.1007/s002200050574](#).
- [BHP12] Arnaud Brothier, Michael Hartglass, and David Penneys, *Rigid C^* -tensor categories of bimodules over interpolated free group factors*, J. Math. Phys. **53** (2012), no. 12, 123525 (43 pages), [arXiv:1208.5505](#), [DOI: 10.1063/1.4769178](#).
- [BP14] Stephen Bigelow and David Penneys, *Principal graph stability and the jellyfish algorithm*, Math. Ann. **358** (2014), no. 1-2, 1–24, [arXiv:1208.1564](#) [MR3157990](#) [DOI: 10.1007/s00208-013-0941-2](#).
- [BR97] Ola Bratteli and Derek W. Robinson, *Operator algebras and quantum statistical mechanics. 2*, second ed., Texts and Monographs in Physics, Springer-Verlag, Berlin, 1997, Equilibrium states. Models in quantum statistical mechanics.
- [EG11] David E. Evans and Terry Gannon, *The exoticness and realisability of twisted Haagerup-Izumi modular data*, Comm. Math. Phys. **307** (2011), no. 2, 463–512, [arXiv:1006.1326](#) [MR2837122](#) [DOI: 10.1007/s00220-011-1329-3](#).
- [EG14] ———, *Near-group fusion categories and their doubles*, Adv. Math. **255** (2014), 586–640, [arXiv:1208.1500](#) [MR3167494](#) [DOI: 10.1016/j.aim.2013.12.014](#).

- [Haa94] Uffe Haagerup, *Principal graphs of subfactors in the index range* $4 < [M : N] < 3 + \sqrt{2}$, *Subfactors* (Kyuzeso, 1993), World Sci. Publ., River Edge, NJ, 1994, [MR1317352](#), pp. 1–38.
- [IMP13] Masaki Izumi, Scott Morrison, and David Penneys, *Quotients of $A_2 * T_2$* , 2013, [DOI : 10.4153/CJM-2015-017-4](#), extended version available as “Fusion categories between $\mathcal{C} \boxtimes \mathcal{D}$ and $\mathcal{C} * \mathcal{D}$ ” at [arXiv:1308.5723](#).
- [Izu01] Masaki Izumi, *The structure of sectors associated with Longo-Rehren inclusions. II. Examples*, *Rev. Math. Phys.* **13** (2001), no. 5, 603–674, [MR1832764](#) [DOI : 10.1142/S0129055X01000818](#).
- [Jon99] Vaughan F. R. Jones, *Planar algebras, I*, 1999, [arXiv:math.QA/9909027](#).
- [Jon01] Vaughan F. R. Jones, *The annular structure of subfactors*, *Essays on geometry and related topics*, Vol. 1, 2, *Monogr. Enseign. Math.*, vol. 38, Enseignement Math., Geneva, 2001, [MR1929335](#), pp. 401–463.
- [Jon12] ———, *Quadratic tangles in planar algebras*, *Duke Math. J.* **161** (2012), no. 12, 2257–2295, [arXiv:1007.1158](#) [MR2972458](#) [DOI : 10.1215/00127094-1723608](#).
- [JP11] Vaughan F. R. Jones and David Penneys, *The embedding theorem for finite depth subfactor planar algebras*, *Quantum Topol.* **2** (2011), no. 3, 301–337, [arXiv:1007.3173](#) [MR2812459](#) [DOI : 10.4171/QT/23](#).
- [Liu13] Zhengwei Liu, *Exchange relation planar algebras of small rank*, 2013, [arXiv:1308.5656](#).
- [Liu15] ———, *Singly generated planar algebras of small dimension, part IV*, 2015, [arXiv:1507.06030](#).
- [LMP15] Zhengwei Liu, Scott Morrison, and David Penneys, *1-supertransitive subfactors with index at most $6\frac{1}{5}$* , *Comm. Math. Phys.* **334** (2015), no. 2, 889–922, [arXiv:1310.8566](#) [MR3306607](#) [DOI : 10.1007/s00220-014-2160-4](#).
- [MP] Scott Morrison and David Penneys, *The affine A and D planar algebras*, In preparation.
- [MP14] Scott Morrison and Emily Peters, *The little desert? Some subfactors with index in the interval $(5, 3 + \sqrt{5})$* , *Internat. J. Math.* **25** (2014), no. 8, 1450080 (51 pages), [arXiv:1205.2742](#) [MR3254427](#) [DOI : 10.1142/S0129167X14500803](#).
- [MP15] Scott Morrison and David Penneys, *Constructing spoke subfactors using the jellyfish algorithm*, *Trans. Amer. Math. Soc.* **367** (2015), no. 5, 3257–3298, [arXiv:1208.3637](#) [MR3314808](#) [DOI : 10.1090/S0002-9947-2014-06109-6](#).
- [MPS15] Scott Morrison, Emily Peters, and Noah Snyder, *Categories generated by a trivalent vertex*, 2015, [arXiv:1501.06869](#).
- [MS12] Scott Morrison and Noah Snyder, *Subfactors of index less than 5, Part 1: The principal graph odometer*, *Comm. Math. Phys.* **312** (2012), no. 1, 1–35, [arXiv:1007.1730](#) [MR2914056](#) [DOI : 10.1007/s00220-012-1426-y](#).
- [Ocn88] Adrian Ocneanu, *Quantized groups, string algebras and Galois theory for algebras*, *Operator algebras and applications*, Vol. 2, *London Math. Soc. Lecture Note Ser.*, vol. 136, Cambridge Univ. Press, Cambridge, 1988, [MR996454](#), pp. 119–172.
- [OP78] Dorte Olesen and Gert Kjaergård Pedersen, *Some C^* -dynamical systems with a single KMS state*, *Math. Scand.* **42** (1978), no. 1, 111–118, [MR500150](#).
- [Pet10] Emily Peters, *A planar algebra construction of the Haagerup subfactor*, *Internat. J. Math.* **21** (2010), no. 8, 987–1045, [arXiv:0902.1294](#) [MR2679382](#) [DOI : 10.1142/S0129167X10006380](#).
- [PP13] David Penneys and Emily Peters, *Calculating two-strand jellyfish relations*, 2013, [arXiv:1308.5197](#), to appear *Pacific J. Math.*
- [TY98] Daisuke Tambara and Shigeru Yamagami, *Tensor categories with fusion rules of self-duality for finite abelian groups*, *J. Algebra* **209** (1998), no. 2, 692–707, [MR1659954](#).