

Rotating Drops Trapped Between Parallel Planes

MARIA ATHANASSENAS

University of Melbourne

November 1993

ABSTRACT. We derive the existence of local minimizers of the functional $\mathcal{F}_\Omega(E)$ describing the energy of a liquid drop $E \subset \mathbb{R}^3$, trapped between two parallel hyperplanes and rotating with constant angular velocity $\sqrt{2\Omega}$, for small $\Omega > 0$.

The study of rotating drops is motivated by problems in astrophysics and physical chemistry. Many physicists and mathematicians have worked on related problems, including Newton, MacLaurin [23], Jacobi [20], Plateau [25], Poincaré [26], Darwin [12], Lord Rayleigh [27], Hölder [19], Appell [2], Lichtenstein [21], Lyttleton [22], Chandrasekhar [8,9], Auchmuty [5], Caffarelli and Friedman [7], Friedman and Turkington [17,18], Brown and Scriven [6].

The question investigated here will be the existence of rotating drops with free boundaries; precisely, of a drop situated between two parallel planes and rotating with constant angular velocity. Of particular interest is the stability of connected drops.

The methods used in this paper will be those introduced by De Giorgi [14,15] for the treatment of variational problems (compare also [16,24]), related to the notion of sets of finite perimeter.

A Lebesgue measurable set $E \subset \mathbb{R}^n$, with characteristic function χ_E , is said to have *finite perimeter* in A , $A \subset \mathbb{R}^n$ open, if the total variation of the vector valued measure $D\chi_E$ satisfies

$$\int_A |D\chi_E| = \sup \left\{ \int_A \chi_E \operatorname{div} g(x) \, dx : g \in C_0^1(A, \mathbb{R}^n), |g(x)| \leq 1 \text{ for } x \in E \right\} < +\infty.$$

We denote by $\Pi_1 = \{x = (y, z) \in \mathbb{R}^2 \times \mathbb{R} : z = 0\}$, $\Pi_2 = \{x = (y, z) \in \mathbb{R}^2 \times \mathbb{R} : z = d\}$, two parallel planes of distance $d > 0$, and by $G = \{x = (y, z) \in \mathbb{R}^2 \times \mathbb{R} : 0 < z < d\}$ the domain between them. The mathematical model of the rotating drop will be to minimize its energy, which is the sum of surface tension, capillarity and rotational energy, and is described by the functional

$$\mathcal{F}_\Omega(E) = \int_G |D\chi_E| + \nu \sum_{i=1}^2 \int_{\Pi_i} \chi_E^\pm d\mathcal{H}^2 - \Omega \int_G |y|^2 \chi_E \, dz \, dy$$

where $\nu, \Omega \in \mathbb{R}$, $0 \leq \nu < 1$, $0 \leq \Omega$. By χ_E^+ we denote the trace of χ_E for $x \in \Pi_i$, $i = 1, 2$, (compare [15,16,24]). The class of admissible sets is chosen to be

$$\mathcal{C} = \left\{ E \subset G \text{ Lebesgue measurable} : \int_G |D\chi_E| < \infty, |E| = 1 \right. \\ \left. \text{and } \int_E y_i dx = 0, i = 1, 2 \right\},$$

that is, the sets E with prescribed volume and barycenter lying on the axis $(0, 0, z)$. The functional \mathcal{F}_Ω describes the energy of a liquid drop rotating at a constant angular velocity $\sqrt{2\Omega}$ around its own barycenter.

As the energy functional is unbounded from below, we shall only treat the question of the existence of local minimizers for \mathcal{F}_Ω .

Let $G(R) = \{(y, z) \in G : |y| < R\}$ for any $R \in \mathbb{R}$. We call $E \in \mathcal{C}$ a *local minimizer* if there exists $R > 0$ such that

- (i) $E \subset\subset G(R)$
- (ii) $\mathcal{F}_\Omega(E) \leq \mathcal{F}_\Omega(F)$ for all $F \in \mathcal{C}$, $F \subset G(R)$.

We define $\mathcal{C}_R = \{E \in \mathcal{C} : E \subset G(R)\}$.

The techniques are the same as those used by Albano and Gonzalez in [1]. In our case, the special difficulty arises from the “free boundary” of E in Π_i , due to the additional capillarity term in the functional.

Related results for rotating drops with obstacles are also obtained by Congedo, Emmer and Gonzalez [11], and Congedo [10] — here the obstacle is assumed to be a graph with a certain growth at infinity. Sturzenhecker [28] treats the cases of pendent rotating drops.

The main result we present is

Main Theorem. *There exists $\Omega_1 > 0$ such that for $0 < \Omega < \Omega_1$, the energy functional \mathcal{F}_Ω has a local minimizer.*

The complete proofs being given in [4], we summarise here the main ideas.

1. General existence results.

Using the standard compactness theorem for $BV(G(R))$ -functions uniformly bounded in the $BV(G(R))$ -norm (see [16]) and the lower semicontinuity of \mathcal{F}_Ω with respect to L^1 -convergence, we obtain the following two results:

Theorem. *Let $R \in \mathbb{R}$, $R > 0$, be such that $|G(R)| > 1$. Then, for each $\Omega \geq 0$ there exists $E_\Omega \in \mathcal{C}_R$ minimizing \mathcal{F}_Ω*

$$\mathcal{F}_\Omega(E_\Omega) = \inf\{\mathcal{F}_\Omega(F) : F \in \mathcal{C}_R\}.$$

Theorem. *For a sequence $\{\Omega_j\}_{j \in \mathbb{N}}$ with $\Omega_j \rightarrow 0$, as $j \rightarrow \infty$, we obtain*

$$E_{\Omega_j} \rightarrow E_0 \quad \text{in } L^1(G(R)),$$

where E_0 is \mathcal{F}_0 -minimizing.

This allows the use of the author’s results in [3], where a detailed discussion of the geometrical properties of E_0 is given.

ROTATING DROPS TRAPPED BETWEEN PARALLEL PLANES

2. A stability result for E_0 .

In the case $\Omega = 0$ the minimizer E_0 is known to be an analytic, rotationally symmetric, periodic surface of constant mean curvature. Furthermore, it intersects Π_i at a constant angle γ , for which $\cos \gamma = \nu$, and consists of at most one period. In \mathbb{R}^3 the possible minimizers are classified: they are the Delaunay surfaces [13]. Using this, we know the possible shapes of drops to which E_{Ω_j} would converge for $\Omega_j \rightarrow 0$. We also prove that for “small” (related to the distance of the planes) volume $|E_0|$, the minimizer cannot be connected and have non-empty intersections with both Π_1 and Π_2 . In this case, E_0 is part of a ball satisfying the volume and boundary (contact angle) conditions. (For more details see [3,4].)

3. Existence of local minimizers.

The proof consists of two steps. First we have:

Theorem. Choose R large enough that $\frac{R}{2} > \max \left\{ \left(\frac{4}{3} \pi \right)^{\frac{1}{3}}, \left(\frac{8}{\pi d} \right)^{\frac{1}{2}} \right\}$. Then there exists $\Omega_0 > 0$ such that, for $0 < \Omega < \Omega_0$, there exists t , $\frac{R}{2} \leq t \leq \frac{3R}{4}$, with

$$\int_{G \cap \{|y|=t\}} \chi_{E_{\Omega}} d\mathcal{H}^2 = 0.$$

Intuitively, the drops E_{Ω_j} concentrate more and more in a neighbourhood of E_0 , given the L^1 -convergence for $\Omega_j \rightarrow 0$. Eventually, there will be some cylinder $\{x \in G : |y| = t\}$, which intersects E_{Ω_j} in at most a set of lower-dimension, for $0 < \Omega < \Omega_0$.

The final step is to show that E_{Ω} has no component outside this cylinder.

Theorem. Choose R as large as above. Then there exists $\Omega_1 > 0$ such that, for $0 < \Omega < \Omega_1$ there exists t , $\frac{R}{2} \leq t \leq \frac{3R}{4}$, with

$$\int_{G(R) \setminus G(t)} \chi_{E_{\Omega}}(x) dx = 0.$$

The idea is to cut off any part outside $G(t)$, rescale what is left axially so as to restore the prescribed volume, and translate so that the barycentre once again lies on the z -axis. The resulting set is then shown to have lower energy. This completes the proof of the main theorem.

REFERENCES

1. ALBANO, S., GONZALEZ, E.H.A., *Rotating drops*, Indiana Univ. Math. Journal **32** (1983), 687–702.
2. APPELL, P.E., *Traité de mécanique rationnelle*, Vol. 4, Chapter 9, Gauthier-Villars, Paris, 1932.
3. ATHANASSENAS, M., *A free boundary problem for capillary surfaces*, Manuscripta Math. **76** (1992), 5–19.
4. ATHANASSENAS, M., *Rotating drops trapped between parallel planes*, (Submitted to Manuscripta Math.).
5. AUCHMUTY, J.E.G., *Existence of axisymmetric equilibrium figures*, Arch. Rational Mech. Anal. **65** (1977), 249–261.
6. BROWN, R.A., SCRIVEN, L.E., *The shapes and stability of captive rotating drops*, Phil. Trans. Roy. Soc. **297** (1980), 51.
7. CAFFARELLI, L.A., FRIEDMAN, A., *The shape of axisymmetric rotating fluid*, J. Funct. Anal. **35** (1980), 109–142.
8. CHANDRASEKHAR, S., *The stability of a rotating liquid drop*, Proc. Roy. Soc. London, Ser. A,

MARIA ATHANASSENAS

9. CHANDRASEKHAR, S., *Ellipsoidal figures of equilibrium*, Yale University Press, New Haven, 1969.
10. CONGEDO, G., *Rotating drops in a vessel. Existence of local minima*, Rend. Sem. Mat. Univ. Padova **72** (1984), 135–156.
11. CONGEDO, G., EMMER, M., GONZALEZ, E.H.A., *Rotating drops in a vessel*, Rend. Sem. Mat. Univ. Padova **70** (1983), 167–186.
12. DARWIN, G.C., *On Jacobi's figure of equilibrium for a rotating mass of fluid*, Proc. Roy. Soc. London **41** (1886), 319–342.
13. DELAUNAY, CH., *Sur la surface de révolution dont la courbure moyenne est constante*, Journ. de mathématiques pures et appliquées **6** (1841), 309–320.
14. DE GIORGI, E., *Sulla proprietà isoperimetrica dell'ipersfera, nella classi degli insiemi aventi frontiera orientata di misura finita*, Atti Accademia Nazionale dei Lincei, Serie VIII, Vol. V, 1958.
15. DE GIORGI, E., COLOMBINI, F., PICCININI, L., *Frontiere orientate di misura minima e questione collegate*, Editrice Tecnico Scientifica, Pisa, 1972.
16. GIUSTI, E., *Minimal surfaces and functions of bounded variation*, Birkhäuser Verlag.
17. FRIEDMAN, A., TURKINGTON, B., *Asymptotic estimates for an axisymmetric rotating fluid*, J. Funct. Anal. **37** (1980), 136–163.
18. FRIEDMAN, A., TURKINGTON, B., *The oblateness of an axisymmetric rotating fluid*, Indiana Univ. Math. J. **29** (1980), 777–792.
19. HÖLDER, E., *Gleichgewichtsfiguren rotierender Flüssigkeiten mit Oberflächenspannung*, Math. Z. **25** (1926), 188–208.
20. JACOBI, C.G.J., *Über die Figur des Gleichgewichts*, Ann. Physik **33** (1834), 229–238.
21. LICHTENSTEIN, L., *Gleichgewichtsfiguren der rotierenden Flüssigkeiten*, Springer-Verlag, Berlin, 1933.
22. LYTTLETON, R.A., *The stability of rotating liquid masses*, Cambridge University Press, 1953.
23. MAC LAURIN, C., *A treatise on fluxions*, T.W. & T. Ruddimas, Edinburgh, 1742.
24. MASSARI, U., MIRANDA, M., *Minimal surfaces of codimension one*, Notas de Matematica, North Holland.
25. PLATEAU, J.A.F., *Experimental and theoretical researches on the figures of equilibrium of a liquid mass withdrawn from the action of gravity*, Annual Report of the Board of Regents of the Smithsonian Institution, Washington, D.C., 1863, pp. 270–285.
26. POINCARÉ, H., *Sur l'équilibre d'une masse fluide animée d'un mouvement de rotation*, Acta Math. **7**, (1885), 259–302.
27. LORD RAYLEIGH, *The equilibrium of revolving liquid under capillary force*, Phil. Mag. **28**, (1914), 161–170.
28. STURZENHECKER, T., *Existence of Local Minima for Capillarity Problems*, PhD Thesis, University of Bonn, 1991.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF MELBOURNE, PARKVILLE, VICTORIA 3052, AUSTRALIA.