

Integrable 1/2 BPS Nahm pole defects in $\mathcal{N} = 4$ SYM

Charlotte Kristjansen

Niels Bohr Institute

Based on:

A. Chalabi, C.K., C. Su, Phys.Lett. B866 (2025), (ArXiv:2503.22598)
C.K., & K. Zarembo, JHEPo8 (2023) 184 (ArXiv:2305.03649),
JHEP 02 (2025) 179 (ArXiv:2412.01972)

Baxter 2025 Exactly Solved Models and Beyond:
Celebrating the Life and Achievements of Rodney James Baxter
September 10th, 2025

AdS/CFT

Conformal operator, $\mathcal{O}$ $\longleftrightarrow$ String state



Eigenstates of integrable super spin chain: $|\mathbf{u}\rangle$

Minahan.
Zarembo '02

Main examples: $\mathcal{N} = 4$ SYM (4D), ABJM theory (3D), ...
Planar limit

AdS/dCFT

Co-dimension d defect $\longleftrightarrow$ Probe brane



(Integrable) boundary state $|\Psi_0\rangle$ of spin chain

De Leeuw, C.K.
Zarembo '15

$\langle\Psi_0|\mathbf{u}\rangle$ is the one-point function $\langle\mathcal{O}\rangle$

Motivation

- Insights on the interplay between conformal symmetry, supersymmetry and integrability
- Tests of AdS/dCFT dictionary for set-ups with and without supersymmetry (so far all positive)
- Exact results for novel types of observables such as one-point functions, bulk-to-boundary correlators etc.
- Interesting connections to statistical physics: matrix product states and quantum quenches.
- Possible cross-fertilization with the boundary conformal bootstrap program.

Plan of the talk

- I. $\frac{1}{2}$ BPS Nahm pole defects in $N=4$ SYM
- II. Integrability properties (in particular of GW-surface defects)
- III. Prediction of wrapping (= finite size) corrections from localization
- IV. Summary & Outlook

1/2 BPS Nahm pole defects in $\mathcal{N} = 4$ SYM

Nahm pole defects :
of co-dimension d
(All 1/2 BPS)

$$\langle \Phi \rangle \sim \frac{\Phi^{\text{cl}}}{r} \quad \longleftarrow \text{distance to defect}$$

- $d = 1$: Domain wall defect:

$$\langle \phi_i \rangle \neq 0, \quad i = 1, 2, 3.$$

Karch &
Randall '02
de Leeuw, CK.
& Zarembo '15

- $d = 2$: Gukov-Witten surface defect:

$$\langle \phi_i \rangle \neq 0, \quad i = 1, 2, \quad \langle A_3 \rangle \neq 0.$$

Gukov & Witten '08
Drukker, Gomis & Matsuura '08

Choi, Gomis & Garcia '24
De Leeuw & Holguin '24
Holguin '25

Chalabi, C.K & Su '25

- $d = 3$ 't Hooft line (monopole):

$$\langle \phi_i \rangle \neq 0, \quad i = 1, \quad \langle A_i \rangle \neq 0, \quad i = 1, 2, 3.$$

Kapustin '05
C.K & Zarembo '23, '24

One-point functions

$$\langle \mathcal{O}_{\Delta}^{\text{bulk}}(x) \rangle = \frac{C}{|x_{\perp}|^{\Delta}}$$

Due to vevs scalar operators can have non-zero 1-pt fcts at tree-level

$$\langle \mathcal{O}_{\Delta}(x) \rangle = \underbrace{(\text{Tr} (\Phi_{i_1} \dots \Phi_{i_L}) + \dots)}_{\sim |s_{i_1} \dots s_{i_L} \rangle} |_{\Phi_i \rightarrow \Phi_i^{\text{cl}}}$$

$$\langle \mathcal{O}_{\Delta}(x) \rangle = \frac{1}{|x_{\perp}|^{\Delta}} \frac{\langle \text{MPS} | \mathbf{u} \rangle}{\langle \mathbf{u} | \mathbf{u} \rangle^{1/2}}, \quad \langle \text{MPS} | = \sum_{\{s_i\}} \text{Tr}(\Phi_{i_1}^{\text{cl}} \dots \Phi_{i_L}^{\text{cl}}) \langle s_{i_1} \dots s_{i_L} |$$

De Leeuw, C.K.
Zarembo '15

1/2 BPS Nahm pole defects in $\mathcal{N} = 4$ SYM

	Domain wall	GW surface		't Hooft line
		Non-rigid	rigid	
Parameters	Discrete Magnetic flux	Continuous	Discrete	Discrete Monopole charge
Probe brane	D5	D3	D3	D1
Geometry	$AdS_4 \times S^2$	$AdS_3 \times S^1$	$AdS_3 \times S^1$	AdS_2
Symmetries	$OSp(4^* 4)$	$PSU(1,1 2)^2 \times SO(2)$	$SU(1,1 2)^2$	$OSp(4^* 4)$
Background, Φ^{cl}	Non-Commutative	Commutative	Non-commutative	Commutative

Gukov Witten surface defects in $\mathcal{N} = 4$ SYM

Ordinary case

Gukov & Witten '08 Drukker, Gomis & Matsuura '08

$$\Phi^{\text{cl}} = \frac{1}{\sqrt{2}z} \text{diag}((\beta_1 + i\gamma_1)\mathbb{1}_{N_1}, \dots, (\beta_M + i\gamma_M)\mathbb{1}_{N_M}), \quad z = re^{i\psi}$$

$$A^{\text{cl}} = \text{diag}(\alpha_1 \mathbb{1}_{N_1}, \alpha_2 \mathbb{1}_{N_2}, \dots, \alpha_M \mathbb{1}_{N_M}) d\psi, \quad \alpha_i \text{ real and periodic}$$

Transverse coordinates: r, ψ

Rigid case

Gukov & Witten '08

$$A^{\text{cl}} = \frac{t_3}{\log \frac{r}{r_0}} d\psi, \quad \Phi^{\text{cl}} = \frac{t_1 + it_2}{\sqrt{2}z \log \frac{r}{r_0}},$$

t_1, t_2, t_3 k -dimensional irreducible representation of $SU(2)$

Corresponds to $\beta, \gamma \rightarrow 0$

Probe brane description

$AdS_5 \times S^5$ background

$$ds^2 = \frac{1}{y^2} (dy^2 + dx_0^2 + dx_1^2 + dr^2 + r^2 d\psi^2) + d\Omega_5$$

$$d\Omega_5 = \cos^2 \theta d\Omega_3 + d\theta^2 + \sin^2 \theta d\phi^2$$

World volume coordinates of D3-brane: $x_0, x_1, r, \psi, (AdS_3 \times S^1)$

Embedding functions:

$$\theta = \frac{\pi}{2}, \quad \phi = \phi(\psi) = \phi_0 - \psi, \quad y = y(r) = \frac{r}{\kappa}$$

Integration constants related to β and γ as

$$\beta + i\gamma = \frac{\sqrt{\lambda}}{2\pi} \kappa e^{i\phi_0}$$

Rigid case: $\beta, \gamma \rightarrow 0$, i.e. $\kappa \rightarrow 0$, cone in AdS_5 shrinks

Tree-level integrability properties of GW defects

Chalabi, CK
& Su '25

	Ordinary	Rigid
$SU(2)$ -sector	Trivial	Integrable $\forall k \in \mathbb{N}, k \geq 2$ *
$SO(6)$ sector	Integrable $\forall \beta_i, \gamma_i$	Integrable for $k = 2$
$SL(2)$ -sector	Non-integrable *	Integrable for $k = 2$

* Also noted in Holguin '25

Integrability test: $Q_{2n+1}|B\rangle = 0, \quad \forall n \in \mathbb{N}$ Pozsgay '17


Overlap formula: KT-relation & recursive strategy Gombor '24

Tree-level one-point functions for rigid GW-defects

For $SO(6)$ -sector, $k = 2$

Chalabi, C.K &
Su '25

$$\langle \mathcal{O}_L(x) \rangle \propto \frac{\langle \text{MPS} | \mathbf{u} \rangle}{\langle \mathbf{u} | \mathbf{u} \rangle^{\frac{1}{2}}} = \sqrt{\frac{Q_2(i/2) \det G_+}{Q_2(0) \det G_-}}$$




NB: Baxter polynomials

NB: A-chiral overlap
 $SO(2) \times SO(4)$ symmetry

Domain wall case

$$\langle \mathcal{O}_L(x) \rangle \propto \sqrt{\frac{Q_1(i/2)Q_2(i/2)Q_3(i/2) \det G_+}{Q_1(0)Q_2(0)Q_3(0) \det G_-}}$$



NB: Baxter polynomials

Chiral overlap
 $SO(3) \times SO(3)$ symmetry

Higher loop integrability

- Domain wall defect:
 - String boundary conditions integrable incl. vevs. Linardopoulos & Zarembo '19
 - Asymptotic all loop formula derived Komatsu & Wang '20 Gombor & Bajnok '20
- Gukov-Witten surface defect:
 - String boundary conditions integrable without vevs Dekel & Oz '11
 - Only hope for rigid case $k = 2$ from field theory Chalabi, C.K. & Su '25
- 't Hooft line:
 - String boundary conditions integrable without vevs Dekel & Oz '11
 - Asymptotic all loop formula up to reflection phase exists Gombor & Bajnok '23

Missing: Finite size (= wrapping) corrections
(TBA, QSC, NLIE,...)

Wrapping corrections from localization

Example: 't Hooft loop (TL)

Consider: $\langle \mathcal{O}_L \rangle_T \equiv \frac{\langle T(C) \mathcal{O}_L(x) \rangle}{\langle T(C) \rangle} = \frac{C_L}{(2|x_\perp|)^L}$

with $\mathcal{O}_L = \frac{1}{\sqrt{L}} \left(\frac{4\pi^2}{\lambda} \right)^{\frac{L}{2}} \text{tr} Z^L$

Localization results for Wilson loop $\langle \mathcal{O}_L \rangle_W$

Okuyama &
Semenoff '07

gives prediction for $\langle \mathcal{O}_L \rangle_T$ via S-duality

Kristjansen
& Zarembo '23

Large-N perturbative prediction for $\langle \mathcal{O}_L \rangle_T$

$$\langle \mathcal{O}_L \rangle_T = \frac{1}{(2r)^L} \frac{1}{\sqrt{L}} \left[\left(\sqrt{1 + \frac{\pi^2}{\lambda}} + \frac{\pi}{\sqrt{\lambda}} \right)^L - \left(\sqrt{1 + \frac{\pi^2}{\lambda}} - \frac{\pi}{\sqrt{\lambda}} \right)^L \right] \equiv \frac{1}{(2r)^L} \mathcal{C}_L$$

In Zhukovski variables: $x + \frac{1}{x} = 2u$, $u = \frac{\pi i}{\sqrt{\lambda}}$

$$\mathcal{C}_L = \frac{1}{i^L \sqrt{L}} \left[x^L + \frac{(-1)^{L+1}}{x^L} \right] = \text{regular} + \text{wrapping}$$

$\sim \lambda^{-L/2} + \dots \quad \sim \lambda^{+L/2} + \dots$

Predicted structure of higher loop corrections

$$c_2 = \frac{1}{\sqrt{2}} \left(\frac{2\pi}{\sqrt{\lambda}} \right)^2 \left(1 + \frac{\lambda}{2\pi^2} - \frac{\lambda^2}{8\pi^4} + \frac{\lambda^3}{16\pi^6} + \dots \right),$$

Kristjansen &
Zarembo '25

$$c_3 = \frac{1}{\sqrt{3}} \left(\frac{2\pi}{\sqrt{\lambda}} \right)^3 \left(1 + \frac{3\lambda}{4\pi^2} \right),$$

$$c_4 = \frac{1}{2} \left(\frac{2\pi}{\sqrt{\lambda}} \right)^4 \left(1 + \frac{\lambda}{\pi^2} + \frac{\lambda^2}{8\pi^4} - \frac{\lambda^4}{128\pi^8} + \dots \right),$$

$$c_5 = \frac{1}{\sqrt{5}} \left(\frac{2\pi}{\sqrt{\lambda}} \right)^5 \left(1 + \frac{5\lambda}{4\pi^2} + \frac{5\lambda^2}{16\pi^4} \right),$$

$$c_6 = \frac{1}{\sqrt{6}} \left(\frac{2\pi}{\sqrt{\lambda}} \right)^6 \left(1 + \frac{3\lambda}{2\pi^2} + \frac{9\lambda^2}{16\pi^4} + \frac{\lambda^3}{32\pi^6} - \frac{\lambda^6}{2048\pi^{12}} + \dots \right).$$

Notice:

- The perturbative series truncates for odd L
- For even L wrapping corrections set in from $\mathcal{O}(\lambda^{2L})$
- The non-wrapping part truncates for all L
- For domain walls: Same structure except $c_{2n+1} = 0$.
- For Gukov-Witten defects truncation without wrapping.

Choi, Gomis,
Garcia '24

Understanding the perturbative series

Closed form of non-wrapping (nw) contributions

$$\mathcal{C}_L^{nw} = \frac{1}{\sqrt{L}} \left(\frac{2\pi}{\sqrt{\lambda}} \right)^L \left\{ 1 + \sum_{n=1}^{[L/2]} \binom{L-n}{n} \frac{L}{L-n} \left(\frac{\lambda}{4\pi^2} \right)^n \right\}$$

Wish to recover this from
quantizing around the monopole background

NB: Same structure for the domain wall defect

$$\mathcal{C}_L^{nw} = \frac{1}{\sqrt{L}} \left(\frac{2\pi}{\sqrt{\lambda}} \right)^L \left\{ 1 + \sum_{n=1}^{[L/2]} \binom{L-n}{n} \frac{L}{L-n} \frac{2B_{L-2n+1} \left(\frac{1+k}{2} \right)}{L-2n+1} \left(\frac{\lambda}{4\pi^2} \right)^n \right\}$$

de Leeuw, Ipsen, Komatsu &
C.K. & Wilhelm '17 Wang '20

Future goal:

Understand the wrapping corrections in the same detail
and compute one-point functions of non-protected operators
exactly using integrability

Quantizing around the monopole background

1. Expand around classical fields $(\Phi_1^{\text{cl}}, A_\mu^{\text{cl}})$

$$A_\mu, \Phi_i, \Psi = \begin{array}{c} \begin{array}{cc} 1 & N-1 \end{array} \\ \left[\begin{array}{c|ccc} \alpha & \beta & \beta & \beta \\ \beta & \gamma & \gamma & \gamma \\ \beta & \gamma & \gamma & \gamma \\ \beta & \gamma & \gamma & \gamma \end{array} \right] \begin{array}{c} 1 \\ N-1 \end{array} \end{array}$$

Due to vevs, non-trivial mixing problem for β components (for simplest set-up).

2. Gauge fix

3. Invert the quadratic part of the action (determine propagators)

Spectral decomposition

$$S = \Phi G^{-1} \Phi$$

$$G(x, y) = \sum_k \Psi_k(x) \frac{1}{\lambda_k} \Psi_k^\dagger(y), \quad G^{-1} \Psi_k(x) = \lambda_k \Psi_k(x)$$

Quantum mechanical problem (field components β)

Quantum mechanical problems involved

I. For $\Phi_2, \Phi_3, \dots, \Phi_6, A_0$ and ghost c

Scalar particle in monopole potential

Dirac '31

II. For $\Phi_1, \vec{A}$:

Scalar coupled to spin-1 particle in monopole potential

Spin-1 particle & monopole
Olsen, Osland & Wu '90

III. For Ψ_α^I ($\alpha, I \in \{1, 2, 3, 4\}$)


Fermions in (non-standard) monopole potential

Standard case
Kazama, Yang &
Goldhaber '77

All are beautiful, exactly solvable quantum mechanical systems

Origin of non-wrapping part of corrections

$$\mathcal{C}_L^{nw} = \frac{1}{\sqrt{L}} \left(\frac{2\pi}{\sqrt{\lambda}} \right)^L \left\{ 1 + \sum_{n=1}^{[L/2]} \binom{L-n}{n} \frac{L}{L-n} \left(\frac{\lambda}{4\pi^2} \right)^n \right\}$$




$$\binom{L-n}{n} \frac{L}{L-n} = \binom{L-n}{n} + \binom{L-2-(n-1)}{n-1}$$



Only contributions from flower diagrams
with varying numbers of leaves

À priori expect contributions from
numerous diagrams with internal vertices



- Same phenomenon observed for the domain wall defect.
- Same phenomenon for ordinary Gukov Witten defects but no wrapping.

Summary & Outlook

- (At least) three examples of dCFT's (co-dimension 1,2,3) amenable to localization, integrability and potentially bootstrap
- Quantization completed for all three cases
- Localization performed, implies intriguing structure of perturbative exp.
- One-pt fcts of non-protected operators obtained from integrability up to wrapping interactions for best understood case ($d = 1$)
- Scene is set for full solution by integrability incl. wrapping
- Boundary bootstrap still to be implemented for non-vanishing vevs.